

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/54, 15/55, 15/60, 15/82, 5/10,</b> <b>A01H 5/00, 5/10</b>	<b>A2</b>	<b>(11) International Publication Number: WO 00/03022</b> <b>(43) International Publication Date: 20 January 2000 (20.01.00)</b>
<b>(21) International Application Number:</b> PCT/US99/15472 <b>(22) International Filing Date:</b> 10 July 1999 (10.07.99)  <b>(30) Priority Data:</b> 09/113,257 10 July 1998 (10.07.98) US  <b>(71) Applicant:</b> CALGENE LLC [US/US]; 1920 Fifth Street, Davis, CA 95616 (US).  <b>(72) Inventors:</b> HAJDUKIEWICZ, Peter, 15821 Timbervalley Road, Chesterfield, MO 60317 (US). MCBRIDE, Kevin, E.; 1309 Marina Circle, Davis, CA 95616 (US). NEHRA, Narender, 14971 Appalachian Trail, Chesterfield, MO 63017 (US). SCHAAF, David, J.; 2780 Brentwood Place, Davis, CA 95616 (US). STALKER, David, M.; 870 W. Southwood, Woodland, CA 95695 (US). STAUB, Jeffrey, M.; 819 "J" Foxspring Drive, Chesterfield, MO 63017 (US). YE, Guangning; 2325 Crimson View Court, Ellisville, MO 63011 (US).  <b>(74) Agent:</b> MONSANTO/G.D. SEARLE; Patent Dept. Central, P.O. Box 5110, Chicago, IL 60680-5110 (US).		<b>(81) Designated States:</b> BR, CA, JP, MX, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> EXPRESSION OF HERBICIDE TOLERANCE GENES IN PLANT PLASTIDS  <b>(57) Abstract</b>  Provided are constructs and methods for expressing herbicide tolerance genes in plastids of plant cells. Constructs include the components of a promoter functional in a plant plastid, a DNA sequence which is capable of conferring tolerance in a plant cell to at least one herbicide compound when said DNA sequence is transcribed in plastids of said plant cell and a transcription termination region. Herbicide tolerance is produced by transforming plastids with the constructs of the invention and growing plant cells comprising the transformed plastids under conditions wherein the DNA sequence is transcribed and plant plastids and cells containing the plastids are rendered tolerant to applications of at least one herbicide compound.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## EXPRESSION OF HERBICIDE TOLERANCE GENES IN PLANT PLASTIDS

5

### INTRODUCTION

#### Technical Field

This invention relates to the application of genetic engineering techniques to  
10 plants. Specifically, the invention relates to compositions and methods for enhancing  
expression of proteins in plant plastids.

#### Background

The plastids of higher plants are an attractive target for genetic engineering. Plant  
15 plastids (chloroplasts, amyloplasts, elaioplasts, etioplasts, chromoplasts, etc.) are the major  
biosynthetic centers that, in addition to photosynthesis, are responsible for production of  
industrially important compounds such as amino acids, complex carbohydrates, fatty acids,  
and pigments. Plastids are derived from a common precursor known as a proplastid and  
thus the plastids present in a given plant species all have the same genetic content. Plant  
20 cells contain 500-10,000 copies of a small 120-160 kilobase circular genome, each  
molecule of which has a large (approximately 25kb) inverted repeat. Thus, it is possible to  
engineer plant cells to contain up to 20,000 copies of a particular gene of interest which  
potentially can result in very high levels of foreign gene expression. In addition, plastids  
of most plants are maternally inherited. Consequently, unlike heterologous genes  
25 expressed in the nucleus, heterologous genes expressed in plastids are not pollen  
disseminated, therefore, a trait introduced into a plant plastid will not be transmitted to  
wild-type relatives.

There remains a need for improved regulatory elements for expression of genes in a  
plant plastid. To date, the expression signals used routinely for plastid transgene

expression derive from endogenous plastid genes. The plastid expression signals are typically derived from promoter regions of highly expressed plastid genes such as the promoter regions from the 16S ribosomal RNA operon (*Prrn*), *psbA* gene (*PpsbA*) or the *rbcL* gene (*PrbcL*). The *psbA* and *rbcL* genes are highly transcribed, but their translation is  
5 controlled by tissue-specific and light-regulated factors which limits their usefulness. In the case of *Prrn*, a synthetic ribosome binding site (RBS) patterned after the plastid *rbcL* gene leader has been typically used to direct translation. However, this *Prrn*/RBS is translated inefficiently due to poor ribosome binding.

A totally heterologous expression system has been used to express plastid genes  
10 (USPN 5,576,198, the entirety of which is incorporated herein by reference). This system is a two component system. The first component is a plastid transgene driven by a T7 bacteriophage gene 10 promoter/leader sequence. The second component is a nuclear gene encoding the T7 Polymerase that is targeted to the plastid compartment. The limitation of this system is the need to create nuclear transformed lines that express the T7 Polymerase  
15 in preferred ways.

Plastids of higher plants present an attractive target for genetic engineering. As mentioned above, plastids of higher plants are maternally inherited. This offers an advantage for genetic engineering of plants for tolerance or resistance to natural or chemical conditions, such as herbicide tolerance, as these traits will not be transmitted to  
20 wild-type relatives. In addition, the high level of foreign gene expression is attractive for engineered traits such as the production of pharmaceutically important proteins.

Expression of nucleic acid sequences encoding for enzymes providing for herbicide tolerance as well as pharmaceutical proteins from plant plastid genome offers an attractive alternative to expression from the plant nuclear genome.

25

## SUMMARY OF THE INVENTION

The present invention provides nucleic acid sequences useful in enhancing expression of a wide variety of genes, both eukaryotic and prokaryotic, in plant plastids.

Furthermore, plastid expression constructs are provided which are useful for genetic engineering of plant cells and which provide for enhanced expression of the EPSP synthase proteins or the hGH protein in plant cell plastids. The transformed plastids should be metabolically active plastids, and are preferably maintained at a high copy  
5 number in the plant tissue of interest, most preferably the chloroplasts found in green plant tissues, such as leaves or cotyledons.

The plastid expression constructs for use in this invention generally include a plastid promoter region capable of providing for enhanced expression of a DNA sequence, a DNA sequence encoding an EPSPS protein or hGH, and a transcription termination  
10 region capable of terminating transcription in a plant plastid.

The plastid promoter region of the present invention is preferably linked to a ribosome binding site which provides for enhanced translation of mRNA transcripts in a plant plastid.

The plastid expression construct of this invention is preferably linked to a construct  
15 having a DNA sequence encoding a selectable marker which can be expressed in a plant plastid. Expression of the selectable marker allows the identification of plant cells comprising a plastid expressing the marker.

In a preferred embodiment, vectors for transfer of the construct into a plant cell include means for inserting the expression and selection constructs into the plastid  
20 genome. This preferably comprises regions of homology to the target plastid genome which flank the constructs.

The constructs of the present invention preferably comprises a promoter sequence linked to a ribosome binding site capable of enhancing the translation of mRNA transcripts in the plant plastid. The ribosome binding site is preferably derived from the T7  
25 bacteriophage gene 10 leader sequence.

Of particular interest in the present invention is the high level of expression of nucleic acid sequences in plant plastids. Of particular interest is the high level expression of nucleic acid sequences encoding for enzymes involved in herbicide tolerance and encoding for pharmaceutical proteins.

The constructs of the present invention preferably comprises a DNA sequence encoding for a 5-Enolpyruvylshikimate-3-phosphate synthase (USPN 5,633,435, the entirety of which is incorporated herein by reference), nitrilase, phytoene desaturase, aprotinin or a DNA sequence encoding Human Growth Hormone (USPN 5,424,199, the  
5 entirety of which is incorporated herein by reference).

Plant cell plastids containing the constructs are also contemplated in the invention, as are plants, plant seeds, plant cells or progeny thereof containing plastids comprising the construct.

The present invention also includes methods for enhanced expression of DNA  
10 sequences in plant plastids.

The invention also includes a method for the enhanced expression of an enzyme conferring herbicide tolerance in a plant cell, by expressing the *Agrobacterium tumefaciens* sp stain CP4 EPSPS in plastids of the plant cell.

In addition, the invention also includes a method for the enhanced expression of an  
15 enzyme encoding hGH in plastids of the plant cell.

Thus, the present invention relates to a chimeric gene containing a herbicide tolerance coding sequence or the coding sequence of a pharmaceutical protein, a plant plastid expression vector containing a promoter operably linked to a T7 Bacteriophage Polymerase gene 10 ribosome binding site capable of enhanced expression in a plant  
20 plastid operably linked to a herbicide tolerance or pharmaceutical coding gene, a plant transformation vector having inserted therein a herbicide tolerance or pharmaceutical coding gene expressed from a plastid promoter linked to a T7 Bacteriophage Polymerase gene 10 ribosome binding site, plant cells transformed using such vectors and plants regenerated therefrom which exhibit a substantial degree of expression of nucleic acid  
25 sequences and proteins and methods for producing such plants and such plants.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide sequence of the G10L ribosome binding site.

Figure 2 provides an amino acid sequence encoding for aprotinin.

5      Figure 3 provides the results of RP-HPLC analysis for characterization of hGH protein expressed in the plastid. Peak I (tallest peak) indicates the expected retention time for properly folded, native 22 kDa GP2000.

Figure 4 provides an electrospray ionization mass spectrometry (MS) analysis using a Micromass Q-ToF electrospray time-of-flight mass spectrometer. In particular, a  
10      series of ions corresponding to the specie(s) present in the sample with varying numbers of protons attached is provided. The axes of the spectrum are intensity versus mass-to-charge ratio of the specie(s) present.

Figure 5 provides a graphic representation of the bioactivity of hGH expressed from a plant plastid. The samples represented on the graph are bovine prolactin (bPL),  
15      hGH expressed from *E. coli* (Ala-hGH), and a null transgenic spiked with bovine prolactin (SPFF Null Spike) as positive controls, a null transgenic (SPFF Null) as a negative control, and transgenic samples from a sepharose column (SPFF Sample, SPFF Sample) and a transgenic sample eluted from the sepharose column at pH3.5 (SPFF pH3.5 Eln).

Figure 6 provides the nucleic acid sequence for the Prn/G10L promoter/RBS  
20      hybrid. The Prn promoter contains the consensus plastid -35 and -10 promoter elements (underlined) and the transcription start sites (GC in bold) for the Plastid-Encoded RNA Polymerase (PEP). The gene 10 leader (G10L) contains a perfect plastid ribosome binding site (RBS, nucleotides in bold).

Figure 7 provides the nucleic acid sequence for the Prn/NEP/G10L::14aaGFP  
25      fusion. The NEP promoter region is underlined (A in bold is transcription start site). The NEP promoter region used extends beyond the consensus sequence both upstream and downstream of the promoter. The initial ATG, the initiator methionine is not counted in the 14 amino acids of GFP.

## 30      DETAILED DESCRIPTION OF THE INVENTION

In accordance with the subject invention, plastid expression constructs are provided which generally comprise a promoter functional in a plant plastid, a ribosome binding site

derived from the T7 Bacteriophage Polymerase gene 10 leader, a DNA sequence encoding for a gene of interest, and a transcription termination region capable of terminating transcription in a plant plastid. These elements are provided as operably joined components in the 5' to 3' direction of transcription.

5 Furthermore, the constructs of the present invention may also include a nucleic acid sequence encoding a peptide capable of targeting said DNA sequence encoding a protein to the thylakoid lumen within the chloroplast.

Of particular interest in the present invention is the use of the plastid expression constructs to direct the high level transcription and translation (expression) of nucleic acid  
10 sequences. Such plastid expression constructs find use in directing the high level expression of DNA sequences encoding for enzymes involved in herbicide tolerance or encoding for the production of pharmaceutical proteins.

Of more particular interest in the present invention is the use of the plastid expression constructs to direct the high level translation of transcribed messenger RNA.

15 DNA sequence and biochemical data reveal a similarity of the plastid organelle's transcriptional and translational machineries and initiation signals to those found in prokaryotic systems. In fact, plastid derived promoter sequences have been reported to direct expression of reporter genes in prokaryotic cells. In addition, plastid genes are often organized into polycistronic operons as they are in prokaryotes.

20 Despite the apparent similarities between plastids and prokaryotes, there exist fundamental differences in the methods used to control gene expression in plastids and prokaryotes. As opposed to the transcriptional control mechanisms typically observed in prokaryotes, plastid gene expression is controlled predominantly at the level of translation and mRNA stability by trans-acting nuclear encoded proteins.

25 Translation is a multi-stage process which first involves the binding of messenger RNA (mRNA) to ribosomes. Beginning at the translation start codon, the mRNA codons are read sequentially as the ribosomes move along the mRNA molecule. The specified amino acids are then sequentially added to the growing polypeptide chain to yield the protein or polypeptide encoded in the mRNA.



As mentioned, the first step in the translation process is the binding of the mRNA molecule to the ribosome. The nature of this interaction (i.e. binding) has been only partially elucidated. Analysis of RNase-resistant oligonucleotides isolated from bacterial translation initiation complexes indicate that a RNA fragment approximately 30 to 40  
5 nucleotides in length comprises the initial ribosome binding site (RBS). Thus, a RBS is hereinafter understood to comprise a sequence of mRNA surrounding the translation start codon which is responsible for the binding of the ribosome and for initiation of translation.

Recently, ribosome binding sites have been identified capable of directing translation in a prokaryotes. For example, a ribosome binding site derived from the T7  
10 bacteriophage gene 10 leader, G10L (USPN 5,232,840, the entirety of which is incorporated herein by reference), has been identified which enhances expression of nucleic acid sequences in prokaryotes.

Herbicides such as N-phosphonomethylglycine, halogenated hydroxybenzonitriles, and norflurazon have been the subject of a large amount of investigation.

15 N-phosphonomethylglycine, commonly referred to as glyphosate, inhibits the shikimic acid pathway which leads to the biosynthesis of aromatic compounds including amino acids, plant hormones and vitamins. Specifically, glyphosate curbs the conversion of phosphoenolpyruvic acid (PEP) and 3-phosphoshikimic acid to 5-enolpyruvyl-3-phosphoshikimic acid by inhibiting the enzyme 5-enolpyruvylshikimate-3-phosphate  
20 synthase (hereinafter referred to as EPSP synthase or EPSPS).

Glyphosate tolerant plants have been produced by transformation of various EPSP synthase genes into the nuclear genome of a plant. A gene for EPSP synthase has been cloned from *Agrobacterium tumefaciens* sp strain CP4 (USPN 5,633,435) and confers a high level of glyphosate tolerance in plants. Furthermore, high levels of glyphosate  
25 tolerance has been achieved in a number of crop plants by fusing EPSPS to a chloroplast transit peptide (CTP) for targeted expression in plastids. In addition, variants of the wild-type EPSPS enzyme have been isolated which are glyphosate tolerant as a result of alterations in the EPSPS amino acid coding sequence (Kishore and Shah, *Ann. Rev. Biochem.* (1988) 57:627-663; Shulze et al., *Arch. Microbiol.* (1984) 137:121-123; Kishore

et al., *Fed. Proc.* (1986) 45:1506). These variants typically have a higher  $K_i$  for glyphosate than the wild-type EPSPS enzyme which confers the glyphosate tolerant phenotype, but these variants are also characterized by a high  $K_m$  for PEP which makes the enzyme kinetically less efficient (Kishore and Shah, *Ann. Rev. Biochem.* (1988) 57:627-663; Sost  
5 et al., *FEBS Lett.* (1984) 173: 238-241; Shulze et al., *Arch. Microbiol.* (1984) 137:121-123; Kishore et al., *Fed. Proc.* (1986) 45:1506; Sost and Amrhein, *Arch. Biochem. Biophys.* (1990) 282: 433-436).

In addition to engineering plants for glyphosate tolerance, plants have also been engineered to tolerate other classes of herbicides such as halogenated  
10 hydroxybenzonitriles, and norflurazon using nucleic acid sequences expressed in the nucleus.

Halogenated hydroxybenzonitriles, such as Bromoxynil, are suggested to act herbicidally by inhibiting the quinone-binding protein complex of photosystem II, inhibiting electron transfer (Van Rensen (1982) *Physiol. Plant* 54:515-520, and Sanders  
15 and Pallett (1986) *Pestic. Biochem. Physiol.* 26:116-122). Herbicides such as norflurazon inhibit the production of carotenoids.

Plants which are resistant to Bromoxynil have been produced by expressing DNA sequences encoding for enzymes capable of detoxifying Bromoxynil (nitrilases) in the plant cell nucleus. DNA sequences encoding for such nitrilases have been cloned from  
20 bacteria such as *Klebsiella pneumoniae* and used to construct vectors to direct the expression of the DNA sequence in plant cell nucleus (USPN 4,810,648, the entirety of which is incorporated herein by reference).

Plants which are resistant to Norflurazon have been engineered by expressing nucleic acid sequences which encode for enzymes in the carotenoid biosynthetic pathway  
25 in plant cell nuclei. For example, by expressing a phytoene desaturase from *Erwinia uredovora* provides tolerance to norflurazon.

While plants transformed to express nucleic acid sequences encoding for such enzymes from the nuclear genome have found utility in engineering herbicide tolerant

plants, it would be increasingly beneficial to obtain herbicide tolerant plants via plastidial expression.

In the examples provided herein, DNA sequences encoding for enzymes involved in herbicide tolerance are used in constructs to direct the expression of the sequences from the plant plastid. DNA sequences encoding for 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), bromoxynil nitrilase (Bxn), phytoene desaturase (crtI (Misawa et al, (1993) *Plant Journal* 4:833-840, and (1994) *Plant Jour* 6:481-489), and acetohydroxyacid synthase (AHAS (Sathasiivan *et al.* (1990) *Nucl. Acids Res.* 18:2188-2193)) are used in the expression constructs of the present invention to direct the expression of said herbicide tolerance nucleotide sequences from the plant plastid.

Transplastomic tobacco plants are identified which are homoplasmic for the DNA sequences of interest encoding said herbicide tolerance genes. Homoplasmic plants demonstrate a high level of protein expression from the plastid. Furthermore, homoplasmic plants demonstrate a high level of tolerance for the respective herbicide. For example, as described in more detail in the example below, plants transformed to express EPSPS from the plastid demonstrate a high level of tolerance for the herbicide glyphosate. In addition, homoplasmic tobacco lines expressing nitrilase or phytoene desaturase demonstrate high levels of tolerance for the herbicides bromoxynil and norflurazon, respectively.

An artisan skilled in the art to which the present invention pertains will recognize that additional sequences may be employed to in the plastid expression constructs of the instant invention to produce herbicide tolerant plants. Other nucleic acid sequence which may find use in the plastid expression constructs herbicide tolerant plants include the *bar* gene for tolerance to glufosinate (DeBlock, *et al.* (1987) *EMBO J.* 6:2513-2519).

Furthermore, additional glyphosate tolerance genes may be employed in the constructs of the present invention. Additional glyphosate tolerant EPSPS genes are described in U.S. Patent Number 5,627,061, Padgett *et al.* (1996) *Herbicide Resistant Crops*, Lewis Publishers, 53-85, and in Penaloza-Vazquez, *et al.* (1995) *Plant Cell Reports* 14:482-487, the entireties of which are incorporated herein by reference.

It should be noted that the herbicide tolerance constructs of the present invention may also include sequences encoding genes involved in other stress tolerance genes, for example insect or disease resistance/tolerance genes. As described in more detail in the examples that follow, plastid expression constructs are used to regenerate plants which are resistant to the herbicide Buctril, which also expresses the *Bacillus thuringiensis* cry1Ac protein.

In addition, the plastid expression constructs also find use in directing the production of human biological proteins (pharmaceutical proteins) from the plant plastid. Nucleic acid sequences encoding for the Human Growth Hormone (hGH) are employed in the plastid expression constructs of the present invention. Furthermore, transplastomic tobacco plants containing such constructs demonstrate a high level of expression of hGH. In addition, the hGH protein expressed from the plant plastid exhibits characteristics of proper processing as well as proper protein folding.

Traditional methods of pharmaceutical protein production generally employ prokaryotic or single cell eukaryotic organisms for expression and large scale production systems. For example, production of the human biologic, Human Growth Hormone (USPN 5,424,199), has been achieved in *Bacillus* and *E. coli* cells.

Another example is the production of aprotinin. Traditional methods for the production of aprotinin have employed the expression of aprotinin in bacteria, or more usually, the extraction of the protein from bovine organs or tissues. Thus, there is a need in the art for an alternative approach for the large scale production of such human biologics.

Human Growth Hormone (hGH) participates in much of the regulation of normal human growth and development. This 22,000 dalton pituitary hormone exhibits a multitude of biological effects including linear growth (somatogenesis), lactation, activation of macrophages, insulin-like and diabetogenic effects among others (Chawla, *Ann. Rev. Med.* (1983) 34:519; Edwards, *et al.*, *Science* (1988) 239:769; Thorner *et al.*, *J. Clin. Invest.* (1988) 81:745). Growth deficiency in children leads to dwarfism, which has been successfully treated for more than a decade by exogenous administration of hGH.

hGH is a member of a family of homologous hormones that include placental lactogens, prolactins, and other genetic and species variants or growth hormone (Nicoll, *et al.*, *Endocrine Reviews*(1986) 7:169). hGH is unusual among these in that it exhibits broad species specificity and binds to either the cloned somatogenic (Leung, *et al.*, *Nature* (1987) 33:537) or prolactin receptor (Boutin, *et al.*, *Cell* (1988) 53:69).

Aprotinin (also known as bovine pancreatic trypsin inhibitor, BPTI) is a basic protein present in several bovine organs and tissues, such as the lymph nodes, pancreas, lungs, parotid gland, spleen and liver.

Aprotinin is known to inhibit various serine proteases, including trypsin, chymotrypsin, plasmin and kallikrein, and is used therapeutically in the treatment of acute pancreatitis, various stages of shock syndrome, hyperfibrinolytic hemorrhage and myocardial infarction. In addition, administration of aprotinin in high doses significantly reduces blood loss in connection with cardiac surgery, including cardiopulmonary bypass (Bidstrup, *et al.* (1989) *Cardiovasc Surg.* 44:640-645)

As demonstrated in more detail in the examples that follow, plastid expression constructs are employed to direct the expression of aprotinin and a human growth hormone from the plant plastid.

Other sequences which may find use in the production of human biologics include sequences encoding for insulin or insulin precursors may find use in the expression constructs of the present invention. The skilled artisan will recognize that many nucleotide sequences encoding for human biologics may be employed in the constructs of the present invention to direct their expression from a plant plastid such as those described in Goodman and Gelman (1990) *Pharmacological Basis of Therapeutics*, Pergaman Press, 8<sup>th</sup> Edition, Sections 14 and 15.

In developing the constructs the various fragments comprising the regulatory regions and open reading frame may be subjected to different processing conditions, such as ligation, restriction enzyme digestion, PCR, *in vitro* mutagenesis, linkers and adapters addition, and the like. Thus, nucleotide transitions, transversions, insertions, deletions, or the like, may be performed on the DNA which is employed in the regulatory regions or the

DNA sequences of interest for expression in the plastids. Methods for restriction digests, Klenow blunt end treatments, ligations, and the like are well known to those in the art and are described, for example, by Maniatis *et al.* (in *Molecular cloning: a laboratory manual* (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

5           During the preparation of the constructs, the various fragments of DNA will often be cloned in an appropriate cloning vector, which allows for amplification of the DNA, modification of the DNA or manipulation of the DNA by joining or removing sequences, linkers, or the like. Preferably, the vectors will be capable of replication to at least a relatively high copy number in *E. coli*. A number of vectors are readily available for  
10          cloning, including such vectors as pBR322, vectors of the pUC series, the M13 series vectors, and pBluescript vectors (Stratagene; La Jolla, CA).

          In order to provide a means of selecting the desired plant cells, vectors for plastid transformation typically contain a construct which provides for expression of a selectable marker gene. Marker genes are plant-expressible DNA sequences which express a  
15          polypeptide which resists a natural inhibition by, attenuates, or inactivates a selective substance, *i.e.*, antibiotic, herbicide *etc.*.

          Alternatively, a marker gene may provide some other visibly reactive response, *i.e.*, may cause a distinctive appearance or growth pattern relative to plants or plant cells not expressing the selectable marker gene in the presence of some substance, either as applied  
20          directly to the plant or plant cells or as present in the plant or plant cell growth media.

          In either case, the plants or plant cells containing such selectable marker genes will have a distinctive phenotype for purposes of identification, *i.e.*, they will be distinguishable from non-transformed cells. The characteristic phenotype allows the identification of cells, cell groups, tissues, organs, plant parts or whole plants containing  
25          the construct.

          Detection of the marker phenotype makes possible the selection of cells having a second gene to which the marker gene has been linked. This second gene typically comprises a desirable phenotype which is not readily identifiable in transformed cells, but

which is present when the plant cell or derivative thereof is grown to maturity, even under conditions wherein the selectable marker phenotype itself is not apparent.

The use of such a marker for identification of plant cells containing a plastid construct has been described by Svab *et al.* (1993, *supra*). In the examples provided below, a bacterial *aadA* gene is expressed as the marker under the regulatory control of chloroplast 5' promoter and 3' transcription termination regions, specifically the regulatory regions of the *psbA* gene (described in Staub *et al.*, *EMBO J.*(1993) 12(2):601-606). Numerous additional promoter regions may also be used to drive expression of the selectable marker gene, including various plastid promoters and bacterial promoters which have been shown to function in plant plastids.

Expression of the *aadA* gene confers resistance to spectinomycin and streptomycin, and thus allows for the identification of plant cells expressing this marker. The *aadA* gene product allows for continued growth and greening of cells whose chloroplasts comprise the selectable marker gene product. Cells which do not contain the selectable marker gene product are bleached. Selection for the *aadA* gene marker is thus based on identification of plant cells which are not bleached by the presence of streptomycin, or more preferably spectinomycin, in the plant growth medium.

A number of markers have been developed for use with plant cells, such as resistance to chloramphenicol, the aminoglycoside G418, hygromycin, or the like. Other genes which encode a product involved in chloroplast metabolism may also be used as selectable markers. For example, genes which provide resistance to plant herbicides such as glyphosate, bromoxynil or imidazolinone may find particular use. Such genes have been reported (Stalker *et al.*, *J. Biol. Chem.* (1985) 260:4724-4728 (glyphosate resistant EPSP); Stalker *et al.*, *J. Biol. Chem.* (1985) 263:6310-6314 (bromoxynil resistant nitrilase gene); and Sathasivan *et al.*, *Nucl. Acids Res.* (1990) 18:2188 (AHAS imidazolinone resistance gene)).

Stable transformation of tobacco plastid genomes by particle bombardment is reported (Svab *et al.* (1990), *supra*) and Svab *et al.* (1993), *supra*). The methods described therein may be employed to obtain plants homoplasmic for plastid expression constructs.

Generally, bombarded tissue is cultured for approximately 2 days on a cell division-promoting media, after which the plant tissue is transferred to a selective media containing an inhibitory amount of the particular selective agent, as well as the particular hormones and other substances necessary to obtain regeneration for that particular plant species. Shoots are then subcultured on the same selective media to ensure production and selection of homoplasmic shoots.

Transplastomic tobacco plants are analyzed for a pure population of transformed plastid genomes (homoplasmic lines). Homoplasmy is verified using Southern analysis employing nucleic acid probes spanning a region of the transgene and chloroplast genome (i.e. the insertion region). Transplastomic plants which are heteroplasmic (i.e. contain a mixture of plastid genomes containing and lacking the transgene) are characterized by a hybridization pattern of wild type and transgenic bands. Homoplasmic plants show a hybridization pattern lacking the wild type band.

Alternatively, homoplasmy may be verified using the polymerase chain reaction (PCR). PCR primers are utilized which are targeted to amplify from sequences from the insertion region. For example, a pair of primers may be utilized in a PCR reaction. One primer amplifies from a region in the transgene, while the second primer amplifies from a region proximal to the insertion region towards the insertion region. A second PCR reaction is performed using primers designed to amplify the region of insertion. Transplastomic lines identified as homoplasmic produce the expected size fragment in the first reaction, while they do not produce the predicted size fragment in the second reaction.

Where transformation and regeneration methods have been adapted for a given plant species, either by *Agrobacterium*-mediated transformation, bombardment or some other method, the established techniques may be modified for use in selection and regeneration methods to produce plastid-transformed plants. For example, the methods described herein for tobacco are readily adaptable to other solanaceous species, such as tomato, petunia and potato.

For transformation of soybean, particle bombardment as well as *Agrobacterium*-mediated nuclear transformation and regeneration protocols have been described (Hinchee



*et al.* USPN 5,416,011, and Christou *et al.* USPN 5,015,580). The skilled artisan will recognize that protocols described for soybean transformation may be used

In *Brassica*, *Agrobacterium*-mediated transformation and regeneration protocols generally involve the use of hypocotyl tissue, a non-green tissue which might contain a low plastid content. Thus, for *Brassica*, preferred target tissues would include microspore-derived hypocotyl or cotyledonary tissues (which are green and thus contain numerous plastids) or leaf tissue explants. While the regeneration rates from such tissues may be low, positional effects, such as seen with *Agrobacterium*-mediated transformation, are not expected, thus it would not be necessary to screen numerous successfully transformed plants in order to obtain a desired phenotype.

For cotton, transformation of *Gossypium hirsutum* L. cotyledons by co-cultivation with *Agrobacterium tumefaciens* has been described by Firoozabady *et al.*, *Plant Mol. Bio.* (1987) 10:105-116 and Umbeck *et al.*, *Bio/Technology* (1987) 5:263-266. Again, as for *Brassica*, this tissue may contain insufficient plastid content for chloroplast transformation. Thus, as for *Brassica*, an alternative method for transformation and regeneration of alternative target tissue containing chloroplasts may be desirable, for instance targeting green embryogenic tissue.

Other plant species may be similarly transformed using related techniques. Alternatively, microprojectile bombardment methods, such as described by Klein *et al.* (*Bio/Technology* 10:286-291) may also be used to obtain nuclear transformed plants comprising the viral single subunit RNA polymerase expression constructs described herein. Cotton transformation by particle bombardment is reported in WO 92/15675, published September 17, 1992. Suitable plants for the practice of the present invention include, but are not limited to, soybean, cotton, alfalfa, oil seed rape, flax, tomato, sugar beet, sunflower, potato, tobacco, *maize*, wheat, rice and lettuce.

The vectors for use in plastid transformation preferably include means for providing a stable transfer of the plastid expression construct and selectable marker construct into the plastid genome. This is most conveniently provided by regions of homology to the target plastid genome. The regions of homology flank the construct to be

transferred and provide for transfer to the plastid genome by homologous recombination, via a double crossover into the genome. The complete DNA sequence of the plastid genome of tobacco has been reported (Shinozaki *et al.*, *EMBO J.* (1986) 5:2043-2049). Complete DNA sequences of the plastid genomes from liverwort (Ohyaama *et al.*, *Nature* 5 (1986) 322:572-574) and rice (Hiratsuka *et al.*, *Mol. Gen. Genet.* (1989) 217:185-194), have also been reported.

Where the regions of homology are present in the inverted repeat regions of the plastid genome (known as IRA and IRB), two copies of the transgene are expected per transformed plastid. Where the regions of homology are present outside the inverted repeat regions of the plastid genome, one copy of the transgene is expected per transformed plastid. The regions of homology within the plastid genome are approximately 1kb in size. Smaller regions of homology may also be used, and as little as 100 bp can provide for homologous recombination into the plastid genome. However, the frequency of recombination and thus the frequency of obtaining plants having transformed plastids decreases with decreasing size of the homology regions.

Examples of constructs having regions of homology the plastid genome are described in Svab *et al.* (1990 *supra*), Svab *et al.* (1993 *supra*) and Zoubenko *et al.* (*Nuc Acid Res* (1994) 22(19):3819-3824).

As described in more detail in the examples below, constructs are described which provide for enhanced expression of DNA sequences in plant plastids. Various promoter/ribosome binding site sequences are employed to direct expression in plant plastids. Promoter sequences of the 16S ribosomal RNA operon (*Prn*) are linked to a ribosome binding site (RBS) derived from the T7 bacteriophage gene 10 leader sequence (G10L). DNA sequences expressed under the regulatory control of the *Prn*/G10L sequence show a significantly higher level of protein expression than those levels obtained under the control of other promoter/RBS combinations, while expression of mRNA may or may not be higher in these plants.

In the examples below, nucleic acid sequences encoding CP4 EPSP synthase (USPN 5,633,435) are placed into expression constructs for expression of EPSP synthase

enzyme from the plant plastid. Furthermore, a DNA sequence encoding for hGH (USPN 5,424,199) is also placed into expression construct for the expression of human growth hormone from the plant plastid. The constructs prepared utilize a ribosome binding site designed after the T7 bacteriophage gene 10 leader (G10L) to increase the expression of  
5 the nucleic acid sequences in the plant plastid.

Plastid expression constructs encoding for the expression of EPSPS and hGH are introduced via a chloroplast transformation vector.

Tobacco lines containing the native encoding sequence to the EPSPS enzyme expressed in plastids under the control of the *Prrn*/G10L promoter/ribosome binding site  
10 sequence demonstrate a significantly higher level of protein expression than those levels obtained from EPSPS expressed under the control of the *Prrn*/*rbcL* RBS sequence. However, EPSPS mRNA is expressed at a higher level in plants expressing CP4 EPSPS from the plastid under the control of the *Prrn*/*rbcL*(RBS). These results indicate that translation from transcripts containing the T7 bacteriophage gene 10 ribosome binding site  
15 is more efficient. In addition, protein expression levels of EPSPS obtained from transplastomic tobacco lines expressing EPSPS under the control of the *Prrn*/G10L RBS provide for a high level of glyphosate tolerance.

Furthermore, transplastomic tobacco lines transformed to express hGH under the control of the *Prrn*/G10L promoter/ribosome binding site sequence demonstrate a  
20 significantly higher level of protein expression than those levels obtained from hGH expressed under the control of the *PpsbA* promoter/RBS sequence.

Increases in protein expression levels of at least approximately 200 fold may be obtained from constructs utilizing *Prrn*/G10L ribosome binding site for expression of EPSPS and hGH over the expression levels obtained from other promoter/RBS  
25 combinations for plastid expression. In addition, protein levels obtained from plastid expression constructs utilizing the *Prrn*/G10L promoter/RBS sequence may accumulate 50 to 3500 fold higher levels than from nuclear expression constructs. Thus, inclusion of the G10L ribosome binding site in plastid expression constructs may find use for increasing the levels of protein expression from plant plastids.

Furthermore, the constructs of the present invention may also include sequences to target the expressed protein to a particular suborganellar region, for example, the thylakoid lumen of the chloroplast. For example, as described in the examples below, a nucleotide sequence encoding a peptide from the plastid genome cytochrome *f* targets the expressed  
5 aprotinin protein to the thylakoid membrane. Such targeting of expressed proteins may provide for a compartmentalization of the protein allowing for increased oxidative stability and proper protein folding.

Thus, the constructs and methods of the present invention provide a means for obtaining transplastomic plants with high level tolerance of herbicides. High levels of  
10 tolerance include tolerance of vegetative tissue when amounts of greater than about 16 oz/acre glyphosate are applied, preferably greater than about 32 oz/acre, more preferably greater than about 64 oz/acre, most preferably greater than about 128 oz/acre. Furthermore, high levels of tolerance can also include tolerance of reproductive tissues when amounts of greater than about 16 oz/acre glyphosate are applied, preferably greater  
15 than about 32 oz/acre, most preferably greater than about 64 oz/acre.

The invention now being generally described, it will be more readily understood by reference to the following examples which are included for purposes of illustration only and are not intended to limit the present invention.

## 20 EXAMPLES

### Example 1 Expression Constructs

Constructs and methods for use in transforming the plastids of higher plants are described in Zoubenko *et al.* (*Nuc Acid Res* (1994) 22(19):3819-3824), Svab *et al.* (*Proc.*  
25 *Natl. Acad. Sci.* (1990) 87:8526-8530 and *Proc. Natl. Acad. Sci.* (1993) 90:913-917) and Staub *et al.* (*EMBO J.* (1993) 12:601-606). Constructs and methods for use in transforming plastids of higher plants to express DNA sequences under the control of a nuclearly encoded, plastid targeted T7 polymerase are described in U.S. Patent Number 5,576,198. The complete DNA sequences of the plastid genome of tobacco are reported

by Shinozaki *et al.* (*EMBO J.* (1986) 5:2043-2049). All plastid DNA references in the following description are to the nucleotide number from tobacco.

The complete nucleotide sequence encoding the tobacco cytochrome *f* (*petA*) is described in Bassham *et al.*, (1991) *J Biol Chem* 266:23606-23610 and Konishi *et al.*

5 (1993) *Plant Cell Physiol* 34:1081-1087.

#### 1A. Promoter/Ribosome Binding Site Sequences

The promoter region of the plastid 16S ribosomal RNA operon (*Prrn*) is linked to a synthetic ribosome binding site (RBS) patterned on the plastid *rbcL* gene leader to create the *Prrn/rbcLRBS* fragment. The *Prrn/rbcLRBS* sequence is as described in Svab *et al.*

10 (1993, *supra*) for the *Prrn/rbcL*(S) fragment.

The promoter region of the plastid *psbA* promoter (*PpsbA*) and terminator sequences (*TpsbA*) are described in Staub *et al.* (1993, *EMBO J.*, 12, 601-606).

The *Prrn/G10L* sequence was constructed by annealing two oligonucleotide sequences, T7lead1 and T7lead2 (Table 1), to create the G10L plastid ribosome binding site (Figure 1). The G10L sequence was ligated to the 3' terminus of the *Prrn* promoter sequence as an *EcoRI/NcoI* fragment to create the *Prrn/G10L* sequence.

**Table 1**

---

<b>T7lead1</b>	5'-AAT TGT AGA AAT AAT TTT GTT TAA CTT TAA GAA GGA GAT ATA CC-3'
<b>T7lead2</b>	5'-CAT GGG TAT ATC TCC TTC TTA AAG TTA AAC AAA ATT ATT TCT AC-3'

---

Chimeric genes are preferably inserted into the expression vector to direct their transcription from the *Prrn* promoter. Thus, in the plastid genome, chimeric genes are transcribed from the *Prrn*/RBS promoter, or the *Prrn/G10L* promoter in the plant plastid. The nucleic acid sequence of the *Prrn/G10L* fusion is provided in figure 6.

#### 1B. CP4 EPSPS Plastid Expression Constructs

A plastid expression vector pMON30117 is constructed from a precursor vector pPRV111B (Zoubenko, *et al.* 1994, *supra*, GenBank accession U12813). The vector pMON30117 carries a multiple cloning site for insertion of a passenger gene in a *Prrn/rbcLRBS/Trps16* expression cassette. The *Prrn/rbcLRBS* sequence is cloned into  
5 pPRV111B vector as an *EcoRI/NcoI* fragment, and the terminator region from the plastid *rps16* gene(*Trps16*) is cloned 3' of the *Prrn* promoter as a *HindIII/NcoI* fragment. The *Trps16* fragment comprises the *rps16* gene 3'-regulatory region from nucleotides 5,087 to 4,939 in the tobacco plasmid DNA.

The pPRV111B backbone of the vector pMON30117 contains a marker gene,  
10 *aadA*, for selection on spectinomycin and streptomycin, and *rps 7/12* for the integration, by homologous recombination, of the passenger DNA into *trnV-rps7/12* intergenic region.

A nuclear expression construct, pMON10154, was prepared as a control for integration into plants by *Agrobacterium*-mediated transformation. In this construct, the CP4 native gene is expressed from the constitutive Figwort Mosaic Virus promoter and the  
15 *Petunia* HSP70 leader, and has the E9 terminator. Targeting to plastids is by the chloroplast transit peptide of the *Petunia* EPSPS translationally fused to the N-terminus of the CP4 gene.

The plastid expression construct pMON30118 was prepared by cloning the native CP4 EPSPS gene fused with the N-terminal five (5) amino acids from the plastid *rbcL*  
20 (described in Svab *et al.*, 1993 *supra*) gene as an *NcoI/SmaI* fragment into the multiple cloning site of the vector pMON30117.

The plastid expression construct pMON30123 is essentially the same as pMON30118 with the exception of the deletion of the N-terminal five (5) amino acids from the plastid *rbcL*.

25 The plastid expression construct pMON30130 was created by replacing the native CP4 EPSPS of pMON30123, with a synthetic CP4 gene. This construct also lacks the N-terminal 5 amino acid fusion from the plastid *rbcL* gene.

The plastid expression construct pMON38773 was constructed by replacing the *Prrn*/RBS sequence of pMON30123 with the *Prrn*/G10L promoter sequence described

above. The EPSPS DNA sequence of pMON38773 also lacks the N-terminal 5 amino acid fusion from the plastid *rbcL* gene.

A plastid expression construct, pMON38766 was constructed using the promoter from T7 phage gene 10 (P-T7), including G10L, CP4 (native) gene coding region, and the  
5 terminator sequence from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON38797 was constructed using the promoter from T7 phage gene 10 (P-T7), including G10L, CP4 (synthetic) gene coding region, terminator from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON38798 was constructed using the promoter  
10 of the 16SrDNA operon (Prn), G10L, CP4 (synthetic) gene coding region, terminator from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON38793 was constructed using the promoter of the 16SrDNA operon (Prn), a synthetic ribosome binding site (RBS) patterned from the plastid *rbcL* gene, the glyphosate tolerant *Petunia* EPSP synthase gene (P-EPSPS,  
15 Padgett, *et al.* (1987) *Arch. Biochem. Biophys.* 258:564-573) carrying the mutation Glycine to Alanine at amino acid position 101, terminator from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON38796 was constructed using the promoter of the 16SrDNA operon (Prn), synthetic ribosome binding site (RBS) patterned from the  
20 plastid *rbcL* gene, the glyphosate tolerant *Achromobacter* (strain LBAA) EPSP synthase gene (U.S. Patent Number 5,627,061, the entirety of which is incorporated herein by reference) carrying the mutation Glycine to Alanine at amino acid position 100 (G100A), terminator from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON45204, was constructed using the promoter  
25 of the 16SrDNA operon (Prn) with the G10L, the glyphosate tolerant *Pseudomonas* (strain LBAA) EPSP synthase gene carrying the mutation Glycine to Alanine at amino acid position 100 (G100A), terminator from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON45201, was constructed using the promoter of the 16SrDNA operon (Prn), synthetic ribosome binding site (RBS) patterned from the

plastid *rbcl* gene, wild-type glyphosate tolerant *Bacillus subtilis* *aroE* (EPSPS)(U.S. Patent Number 5,627,061) gene, terminator from plastid *rps16* gene (*Trps16*).

A plastid expression construct, pMON45259, was constructed using the promoter of the 16SrDNA operon (*Prrn*) with the G10L sequence functionally associated with the nucleic acid sequence encoding the synthetic CP4 protein having an additional sequence at the N-terminus encoding the first 14 amino acids of the green fluorescent protein (GFP) (GKGEELFTGVVPSM). The sequence encoding the 14 amino acid GFP fusion begins at the glycine in the second position of the protein. The construct also contains the *rps16* terminator.

Another plastid expression construct, pMON49218, was constructed to express the synthetic CP4 sequence with the 14 amino acid GFP fusion from the promoter region of the 16SrDNA operon having the nuclear-encoded RNA polymerase region (*Prrn*PEP+NEP), and the terminator region from the plastid *rps16* gene. The DNA sequence of the *Prrn*/NEP/G10L::14aaGFP fusion is provided in figure 7.

#### 1C. Bucril (*bxn*) Plastid Expression Constructs

The *bxn* herbicide resistance gene (U.S. Patent Number 4,810,648, the entirety of which is incorporated herein by reference) was removed from the plasmid pBrx47 as an *Nco* I to *Asp*718 restriction fragment and cloned into *Nco*/*Asp*718 cut pUC120 resulting in plasmid pBrx87. Plasmid pBrx87 was then digested with *Nco*/*Xba* and cloned into the *Nco*/*Xba* sites of the plasmid pLAA21 which contains the *Prrn* plastid promoter and the *rpsL* 3' region for plastid expression. The resulting plasmid was designated pBrx89. Plasmid pBrx89 was digested with *Sac* I and *Hind* III and the 1.5 kb chimeric *bxn* gene with plastid expression signals was inserted into the *Sac*/*Hind* III sites of the tobacco plastid homology vector pOVZ44B (Zoubenko et al, Nuc Acids Res 22: 3819-3824 (1994)) to create plasmid pCGN5175.

To construct plasmid pCGN6114, plasmid pBrx90 (a Bluescript plasmid containing the *bxn* gene encoding the bromoxynil specific nitrilase) was digested with *Nco* I/*Asc* I and the *bxn* structural gene was substituted for the GUS gene in the *Nco*/*Asc*



digested plasmid pCGN5063 resulting in plasmid pCGN6107. This plasmid contains the the *bxn* gene under the control of the T7 promoter/gene10 leader at the 5' end and the *psbA*/T7 hybrid transcriptional terminator at the 3' end of the chimeric gene. This T7 promoter/*bxn* chimeric gene was excised from pCGN6107 as a *Hind* III/*Not* I DNA segment and moved into the chloromphenical plasmid BCSK+ (Stratagene) at the *Hind* III/*Not* sites to create plasmid pCGN6109. The chimeric gene was then moved as a *Hind* III/*Not* fragment from pCGN6109 into the chloroplast homology vector pOVZ44B described above to create plasmid pCGN6114. Tobacco plants transformed with pCGN6114 require the T7 RNA polymerase be provided in the plant plastid background to activate transcription of the chimeric *bxn* gene via the T7 promoter. This system has previously been detailed in McBride et al PNAS 91:7301-7305 (1994) and McBride et al US patent number 5,576,198.

#### 1D. BXN/AHAS Plastid Expression Constructs

A plastid expression construct, pCGN5026, is prepared to direct the expression of BXN and AHAS from the plant plastid. The AHAS nucleotide sequence (described in EP Publication Number 0 525 384 A2, the entirety of which is incorporated herein by reference) is translationally linked to the BXN nucleotide sequence (U.S. Patent Number 4,810,648, the entirety of which is incorporated herein by reference). The AHAS structural gene encoding acetohydroxyacid synthase was cloned from the plasmid pCGN4277 as an *Nco* I to *Age* DNA fragment into the *Nco*/*Xma* sites of plasmid pUC120 to create plasmid pCGN5022. This plasmid was then digested with the enzymes *Bam*H I and *Pst* and a 1.3 kb *Bam*/*Pst* DNA segment containing the *bxn* gene encoding the bromoxynil-specific nitrilase was excised from the plasmid pBrx26 and cloned into the *Bam*/*Pst* sites of pCGN5022 to create plasmid pCGN5023. Plasmid pCGN5023 contained a 3.3 kb DNA segment containing the AHAS/*bxn* operon segment and this fragment. This plasmid was cut at the unique *Pst* site and this *Pst* site was removed and replaced with a synthetic linker containing a unique *Xba* I restriction site generating plasmid pCGN5024. Plasmid pCGN5024 was digested with *Nco*/*Xba* and the 3.3 kb *Nco*/*Xba* DNA fragment was

cloned into the plastid promoter cassette vector pLAA21(Pst) that had been digested with *Nco* and *Xba* to remove the GUS gene. The plasmid resulting from this cloning was designated plasmid pCGN5025 and contained the herbicide operon under the control of the plastid promoter *Prm* and the *rpsL* 3' DNA segment. The entire chimeric herbicide  
 5 operon under the control of the plastid expression elements was excised from pCGN5025 as a *Sac* I/*Pst* DNA fragment and cloned into the *Sac*/*Pst* sites of the plastid homology cassette vector pOVZ44B (Zoubenko *et al.*, *Nuc Acids Res* 22:3819-3824 (1994)) to facilitate transfer into the tobacco chloroplast genome.

#### 10 1E. Bt *cryI*Ac and *bxn* Plastid Expression Construct

Plasmid pBrx9 (Stalker and McBride, (1987) *J Bacteriol* 169:955-960), an original clone from *Klebsiella* containing a *bxn* gene DNA segment, was used as a template to generate an ~450 bp *Bam*H I/*Cla* I PCR DNA fragment that encompasses the N-terminal end of the *bxn* gene and includes 44 bp of the 5' untranslated portion of the native gene.  
 15 This fragment was exchanged with the ~400 bp *Bam*/C*la* fragment in the plasmid pBrx90 resulting in plasmid pBrx90.1. This plasmid contains the entire *bxn* gene and the 44 bp untranslated 5' DNA segment. The *bxn* gene was excised from plasmid pBrx90.1 as a *Bam*/A*sc* I DNA segment and inserted into plasmid pCGN5146 at the *Bgl* II/A*sc* I sites to generate plasmid pCGN5191. Plasmid pCGN5146 is a pKK233-2 (Pharmacia) derivative  
 20 containing the full-length *cryI*Ac gene encoding the HD-73 Bt protoxin. Plasmid pCGN5191 then contains the *cryI*Ac and *bxn* genes in an operon configuration with the *bxn* gene being the distal gene in the operon. Both genes are under the control of the *Ptac* promoter for *E coli* expression in 5191. Plasmid pCGN5191 was digested with *Nco*/A*sc* and the *Nco*/A*sc* DNA fragment containing the Bt/*bxn* operon was cloned into the *Nco*/A*sc*  
 25 sites of the chloroplast homology vector pCGN5155, a derivative of pOVZ44B. The resulting plasmid, pCGN5197 contains the Bt/*bxn* operon under the control of the *Prm* plastid promoter and *rpsL* transcription terminator regions. This plasmid facilitated transfer of the Bt/*bxn* chimeric operon into the tobacco plastid genome.

#### 1F. Phytoene desaturase Plastid Expression Constructs

The *crtI* gene was obtained as a *Hind* III/*Sal* I PCR fragment from the original plasmid containing the *Erwinia carotova crt* operon (Misawa et al, (1994) *Plant Jour* 6:481-489)) and cloned as a *Hind* III/*Sal* I DNA segment into BCSK+ (Stratagene) at the  
5 *Hind* III/*Sal* I sites to generate plasmid pCGN5172. The *crtI* fragment was cloned from pCGN5172 as an *Nco* I/*Sal* I fragment into pCGN5038 (a derivative of pOVZ44B) to create the plastid expression construct pCGN5177. This construct directs the expression of the *crtI* sequence from the *Prm* promoter and the *rps16* terminator sequence. This plasmid facilitated the transfer of the chimeric *crtI* gene into the tobacco plastid genome.

10

#### 1G. hGH Expression Constructs for Plant Transformation

##### Nuclear Expression Constructs

The construct pWRG4747 was constructed to direct the expression of hGH in the plant nuclear genome. This vector contains the hGH operably linked to the Figwort  
15 Mosaic Virus promoter (USPN 5,378,619, the entirety is incorporated herein by reference) and the CTP2 leader for directing the hGH protein into the plastid. The FMV/CTP2L::hGH::NpA fragment is cloned along with the DNA sequence conferring resistance to Kanamycin between the right and left borders (RB and LB) of the transfer DNA (tDNA) of *Agrobacterium tumefaciens* to direct the integration into the nuclear  
20 genome.

The nuclear transformation vector pWRG4744 contains essentially the same elements as pWRG4747 except the construct lacks the CTP2 leader and the hGH protein is directed to the plant cell cytoplasm.

##### Plastid Expression Constructs

25 The plastid expression vector pWRG4838 was constructed using the full length hGH gene expressed from the promoter region from the *psbA* gene and the *psbA* gene terminator, *PpsbA* and *TpsbA* respectively (described in Staub *et al.* (1993), *supra*). This chimeric promoter-gene-terminator fusion (*PpsbA*::hGH:: *TpsbA*) is cloned adjacent to the selectable marker gene *aadA* also driven by the plastid expression elements of the *psbA*

gene. The two chimeric gene sequences are cloned into a vector between two sequences which direct the integration of the chimeric gene sequences into the tobacco plastid genome upstream of the plastid 16SrDNA. This is joined to a 1 kb Ampicillin resistance gene which provides for selection of *E. coli* containing the construct and the pUC origin of replication for plasmid maintenance in *E. coli*.

The plastid expression construct pMON38755 was prepared using the hGH DNA sequence translationally fused at the N-terminus with the yeast ubiquitin gene, creating the Ubi-hGH fusion gene. The Ubi-hGH fusion gene is cloned next to the *aadA* gene for selection of transplastomic tobacco on media containing spectinomycin or streptomycin (from pPRV112B described in Zoubenko *et al.* (1994) *supra*). Sequences are included for the homologous recombination of sequences encoding for hGH and *aadA* expression. These sequences are obtained from the vector pPRV112B described in Zoubenko *et al.* (1994, *supra*). This is joined to a 1 kb Ampicillin resistance gene which provides for selection of *E. coli* containing the construct and the pUC origin of replication for plasmid maintenance in *E. coli*.

The plastid expression construct pMON38794 contains essentially the same elements as pMON38755, with the following exception. The 0.15 kb *psbA* promoter sequence is replaced with the *Prrn*/G10L promoter sequence described above.

#### 20 1H. Constructs for the Expression of Aprotinin in Plastids

A series of constructs were prepared to direct the expression of the pharmaceutical protein aprotinin from the plastid. The nucleic acid sequence encoding for aprotinin (Figure 2) was cloned into a plastid expression construct to control the expression of aprotinin from the T7 gene 10 leader promoter which is induced from a nuclearly expressed, plastid targeted T7 Polymerase. The constructs used in which the aprotinin sequence was cloned are as described in U.S. Patent Number 5,576,198, the entirety of which is incorporated herein by reference. The plastid transformation vector pCGN6146 is designed by replacing the DNA sequence encoding for GUS from pCGN4276 (described in USPN 5,576,198) with the coding sequence of aprotinin. The tobacco plastid

transformation construct pCGN6147 contains the same elements as pCGN6146 except pCGN6147 contains the six 5' amino acids of the GUS encoding sequence ligated to the 5' terminus of the aprotinin encoding sequence. The six amino acids of the 5' terminus of the GUS nucleotide sequence are included to aid in the translation of the aprotinin protein.

5 The tobacco plastid transformation vector pCGN6156 is essentially the same as pCGN4276 except the coding region of aprotinin is cloned to the 3' end of the GUS coding sequence. Thus, pCGN6156 contains as operably linked the T7 promoter, a DNA sequence encoding for GUS fused with the DNA sequence encoding for aprotinin and the psbA 3' transcription termination sequence.

10 A plastid expression construct, pCGN6154, was constructed from pCGN4276 by replacing the GUS coding sequence with the aprotinin protein operably linked to the 3' terminus of the coding sequence of cytochrome *f* (*petA*) of the tobacco chloroplast. Thus, pCGN6154 contains the T7 promoter sequence operably linked to the nucleotide sequence of *petA* and aprotinin. The *petA* sequence is included to direct the expressed aprotinin  
15 protein to the thylakoid.

## Example 2 Plant Transformation

### 2A. Nuclear Transformation

Tobacco plants transformed to express the constructs pWRG4744 and pWRG4747  
20 in the nucleus of a plant cell may be obtained as described by Horsch *et al.* (*Science* (1985) 227:1229-1232).

### 2B. Plastid Transformation

Tobacco plastids are transformed by particle gun delivery of microprojectiles as described by Svab and Maliga (*Proc. Natl. Acad. Sci.* (1993) 90:913-917), and described  
25 here.

Dark green, round leaves are cut, preferably from the middle of the shoots, from 3-6 week old *Nicotiana tabacum* cv. Havana which have been maintained *in vitro* on hormone free MS medium (Murashige and Skoog, (1962) *Physiol Plant.* 15, 473-497) supplemented with B5 vitamins in Phytatrays or sundae cups with a 16 hour photoperiod at

24°C. Each cut leaf is then placed adaxial side up on sterile filter paper over tobacco shoot regeneration medium (TSO medium: MS salts, 1mg/l *N*<sup>6</sup>-benzyladenine, 0.1mg/l 1-naphthaleneacetic acid, 1 mg/l thiamine, 100mg/l inositol, 7g/l agar pH 5.8 and 30g/l sucrose). Leaves are preferably placed in the center of the plate with as much contact with the medium as possible. The plates are preferably prepared immediately prior to use, but may be prepared up to a day before transformation by particle bombardment by wrapping in plastic bags and storing at 24°C overnight.

Tungsten or gold particles are sterilized for use as microcarriers in bombardment experiments. Particles (50mg) are sterilized with 1 ml of 100% ethanol, and stored at -20°C or -80°C. Immediately prior to use, particles are sedimented by centrifugation, washed with 2 to 3 washes of 1 ml sterile deionised distilled water, vortexed and centrifuged between each wash. Washed particles are resuspended in 500 µl 50% glycerol.

Sterilized particles are coated with DNA for transformation. Twenty-five microliter aliquots of sterilized particles are added to a 1.5 ml microfuge tube, and 5 µg of DNA of interest is added and mix by tapping. Thirty-five microliters of a freshly prepared solution of 1.8M CaCl<sub>2</sub> and 30 mM spermidine is added to the particle/DNA mixture, mixed gently, and incubated at room temperature for 20 minutes. The coated particles are sedimented by centrifuging briefly. The particles are washed twice by adding 200µl 70% ethanol, mixing gently, and centrifuging briefly. The coated particles are resuspended in 50µl of 100% ethanol and mixed gently. Five to ten microliters of coated particles are used for each bombardment.

Transformation by particle bombardment is carried out using the PDS 1000 Helium gun (Bio Rad, Richmond, CA) using a modified protocol described by the manufacturer.

Plates containing the leaf samples are placed on the second shelf from the bottom of the vacuum chamber and bombarded using the 1100 p.s.i. rupture disk. After bombardment, petriplates containing the leaf samples are wrapped in plastic bags and incubated at 24°C for 48 hours.

After incubation, bombarded leaves are cut into approximately 0.5 cm<sup>2</sup> pieces and placed abaxial side up on TSO medium supplemented with 500 µg/ml spectinomycin.

After 3 to 4 weeks on the selection medium, small, green spectinomycin resistant shoots will appear on the leaf tissue. These shoots will continue to grow on spectinomycin containing medium and are referred to as primary putative transformants.

When the primary putative transformants have developed 2 to 3 leaves, 2 small  
5 pieces (approximately 0.5 cm<sup>2</sup>) are cut from each leaf and used for either selection or for a second round of shoot regeneration. One piece is placed abaxial side up on plates containing TSO medium supplemented with 500 µg/ml spectinomycin, and the other piece is placed abaxial side up on TSO medium supplemented with 500 µg/ml each of spectinomycin and streptomycin. Positive transformants are identified as the shoots which  
10 form green callus on the TSO medium containing spectinomycin and streptomycin.

After 3 to 4 weeks, the tissue placed on TSO medium containing only spectinomycin, which has been identified as positive on the TSO medium with spectinomycin and streptomycin, will develop green shoots. Two to four shoots of each positive transformant are selected and transferred to TSO medium supplemented with 500  
15 µg/ml spectinomycin for generation of roots. Southern analysis is performed on 2 shoots to confirm homoplasmy as described below. Shoots from homoplasmic events are transferred to the greenhouse for seed production, while transformants which are not homoplasmic are sent through a second round of regeneration on TSO medium with 500 µg/ml spectinomycin to attain homoplasmy.

20

### **Example 3 Analysis of Transplastomic Tobacco Plants Transformed with Herbicide Tolerance Constructs**

#### **3A. Southern Analysis**

Transformed plants selected for marker *aadA* marker gene expression are analyzed  
25 to determine whether the entire plastid content of the plant has been transformed (homoplasmic transformants). Typically, following two rounds of shoot formation and spectinomycin selection, approximately 50% of the transgenic plantlets which are analyzed are homoplasmic, as determined by Southern blot analysis of plastid DNA. Homoplasmic plantlets are selected for further cultivation.

Genomic DNA is isolated from transformed tobacco plants, electrophoresed, and transferred to filters as described in Svab *et al.* ((1993), *Proc Natl Acad Sci*, 90:913-917).

Homoplasmic tobacco plants transformed to express CP4 EPSPS in plastids were identified using a probe prepared from a 2.4 kb *EcoRI/EcoRV* fragment from the vector pOVZ2 (similar to pOVZ15 described in Zoubenko, *et al.* 1994, *supra*). The 2.4 kb probe  
5 fragment encompasses part of the targeting sequence.

Results of the Southern hybridizations identified 3 homoplasmic lines from tobacco transformed with the constructs pMON30123 and pMON30130 and 1 line from tobacco transformed with pMON38773 for further analysis.

10 The complete disappearance of the 3.27 Kb native tobacco *Bam*HI fragment in the lines 30123-19-1A, 30123-23-2A, 30123-18-1B, 30130-51-2A, 30130-51-2P, 30130-57-1P, and 38773-6 with a probe covering the region of integration, and the appearance of expected sized bands for the inserted DNA fragments in those transformants, 5.14 kb and 0.9 kb, establishes that the transformed plants are homoplasmic for the intended  
15 constructs.

Results of the Southern hybridizations identified 3 homoplasmic lines from tobacco transformed with pCGN5177, lines 74-1B-P, 74-2 and 74-7.

Transplastomic 5175 and 6114 tobacco lines were analyzed by Southern hybridization for homoplasmy as described above. Results of the Southern hybridizations  
20 identified 4 homoplasmic lines from tobacco transformed with pCGN6114.

Results from hybridizations of 5175 transplastomic tobacco lines identified one line, 76-4A-F, as homoplasmic, and a second line as 95% homoplasmic.

Homoplasmic tobacco plants transformed to express BXN/AHAS in plastids were identified using Southern hybridizations as described above.

25 Results of the Southern hybridizations identified 14 homoplasmic lines from tobacco transformed with pCGN5026. The filters were reprobbed with a BXN gene fragment, and 21 lines were found to contain BXN, 14 lines of which were homoplasmic.

### 3B. Northern Analysis



In order to determine the level of transcription of the EPSPS, BXN or AHAS mRNA expressed in the transplastomic tobacco plants, Northern blot hybridizations were performed with total RNA isolated from each of the lines identified. Total RNA was isolated using TRIzol reagent (Gibco-BRL Life Technologies, Gaithersburg, MD) according to the manufacturers protocol. Total RNA, 2µg, was separated on a denaturing agarose gel and transferred to nylon membrane (Maniatis *et al.*, 1989, *supra*). Radioactive probes for hybridizations were prepared using random primer labeled (using Random Primer labeling kit from Boehringer Mannheim) CP4 EPSPS, phytoene desaturase, BXN, or AHAS fragments and hybridizations were carried out in 2x SSPE (Maniatis, *et al.*, 1989, *supra*), at 60°C. Filters were stripped and reprobed with a plastid 16S ribosomal RNA gene probe (from pPRV112A, Zoubenko, *et al.*, 1994, *supra*) to confirm homogenous loading of RNA on the filter.

Results of the Northern hybridizations performed with EPSPS probes demonstrate that all seven (7) lines examined express CP4 EPSPS mRNA. Hybridizations performed with the 16S ribosome probe confirm that denaturing gels were loaded with similar amounts of total RNA for each sample. Furthermore, transplastomic tobacco lines expressing EPSPS from the *Prrn/rbcL*(RBS) (pMON30123) regulatory elements express EPSPS mRNA to higher levels than tobacco plants homoplasmic for EPSPS controlled by the *Prrn/G10L* (pMON38773) promoter/RBS sequences.

Results of Northern hybridizations performed with BXN, AHAS and *crtI* probes demonstrates that all homoplasmic 5026, 5175, and 5177 tobacco lines expressed *crtI*, BXN and/or AHAS mRNA.

### 3C. Western Blot Analysis of Tobacco CP4 EPSPS

To determine the expression of the EPSPS Western blot analysis was performed on a single line from each construct, pMON30123, pMON30130, and pMON38773.

Total soluble protein was extracted from frozen leaf tissue by grinding 250 mg tissue in 250µl of PBS buffer (1 mM KH<sub>2</sub>PO<sub>4</sub>, Na<sub>2</sub>HPO<sub>4</sub>, 0.137M NaCl, 2.7 mM KCl pH 7.0) containing protease inhibitors. The homogenate is centrifuged for 5 minutes, and the

supernatant is transferred to a fresh tube. The concentration of the protein in the supernatant is determined using a protein concentration assay (BioRad, Richmond, CA).

Extracted total protein is electrophoresed on a 4-20% SDS-PAGE gel (Sigma, St Louis, MO), and transferred to PVDF membrane in 1x SDS-PAGE buffer (Maniatis *et al.* 1989, Cold Spring Harbor Press). Standards of quantitated purified CP4 EPSPS protein were used to quantify the expression of the CP4 EPSPS as expressed in the plant plastid.

Western hybridizations are performed as described in Staub and Maliga (1993) *EMBO Journal*, 12(2) 601-606, except using antibodies raised to EPSPS. PVDF membranes containing the transferred electrophoresed protein were incubated in a blocking solution of PBS buffer containing 0.05% Tween-20 (PBS-T) and 5% milk overnight at 4°C. The membranes are then incubated in a solution of PBS-T containing 1% milk and a primary antibody raised in goats to the CP4 EPSPS for 2 hours at room temperature. The membranes are washed three times in a solution of PBS-T containing 0.1% milk, each wash for 5 minutes at room temperature. The membranes are then incubated in a solution of PBS-T containing 1% milk and sheep anti-goat antibody for 1 hour at room temperature, and washed again in PBS-T containing 0.1% milk, three times for 10 minutes at room temperature. A final wash using only PBS-T is performed before developing the membranes using a nonradioactive detection kit (ECL, Amersham).

20

Table 2

Construct Number	% Total Soluble Protein
pMON30123	0.001
pMON30130	0.002
pMON38773	0.2
pMON38798	0.2
pMON45259	>12.0
pMON49218	>12.0

The results listed in Table 2 demonstrate that significant increases in the level of EPSPS protein may be obtained from plants transformed to express EPSPS from the

25

*Prrm*/G10L promoter. These results demonstrate that EPSPS expression driven by the *Prrm*/rbcLRBS regulatory sequences may produce approximately 0.001% of the total soluble protein as EPSPS, while in plants expressing EPSPS from the *Prrm*/G10L regulatory sequences express 0.2% of the total soluble protein as EPSPS. Subsequent lines have demonstrated total soluble protein of about 1% EPSPS when expressed from the *Prrm*/G10L regulatory sequences. These results, taken together with the results of the Northern hybridizations above, indicate that more efficient translation may be obtained from the G10L ribosome binding site.

Furthermore, plastid expression constructs containing the N-terminal 14 amino acid from GFP demonstrated high levels of protein expression. Transplastomic lines containing either pMON45259 or pMON49218 demonstrated total soluble protein of greater than 12% CP4 EPSPS.

Western immunoblot hybridization were also performed on 2 homoplasmic 5026 tobacco lines as described above, using antibodies raised against bromoxynil. The results of Western immunoblot analysis of total soluble protein extracted from tobacco lines transformed with pCGN5026 demonstrated that both homoplasmic lines produced nitrilase protein.

Western immunoblot analysis was performed as described above from total protein extracted from tobacco lines transformed with pCGN6114 and pCGN5197.

The results of the analysis demonstrated that bromoxynil was produced in 6114 tobacco lines ranging from 1% to 2% of the total soluble leaf protein.

The results of the Western analysis of the 20 5197 tobacco lines demonstrated that bromoxynil and Bt were both produced as 1% of the total soluble leaf protein.

### 25 3D. Analysis of EPSPS Enzyme Activity

The EPSPS enzyme activity in transplastomic tobacco plants containing the plastid expression vector pMON38773 was determined using a high pressure liquid chromatography (HPLC) assay.

Methods for the analysis of EPSPS enzyme activity are described in Padgett *et al.* (*J. Biol. Chem.* (1988)263:1798-1802 and *Arch. Biochem. Biophys.* (1987)258:564-573) and Wibbenmeyer *et al.* (*Biochem. Biophys. Res. Commun.* (1988)153:760-766).

The results are summarized in Table 3 below.

Table 3

<u>Nuclear</u> Enzymatic Activity Range	<u>Nuclear</u> % Total Plants In Range	<u>Chloroplast</u> 38773-6
1-3.7 $\mu\text{mol/mg}$	1%	
>0.1 $\mu\text{mol/mg}$	16%	
>10 nmol/mg	55%	16.39 nmol/mg
>1 nmol/mg	32%	
0 nmol/mg	3%	

These results demonstrate that EPSPS expression in plastids produces active EPSPS enzyme.

5

### 3E. Analysis for Glyphosate Tolerance

A transplastomic tobacco line homoplasmic for the construct pMON38773 was tested *in vitro* to determine the highest level of glyphosate tolerance. Explant tissue was prepared from leaf pieces of nontransgenic wild type tobacco control, *Havanna*, plants and the homoplasmic tobacco line 38773-6 and cultured for regeneration of shoots on TSO medium (described above) supplemented with glyphosate levels of 50 $\mu\text{M}$ , 75 $\mu\text{M}$ , 100 $\mu\text{M}$ , 150 $\mu\text{M}$  and 200 $\mu\text{M}$ . The results are summarized in Table 4 below. The number of explants producing shoots was determined at 3 weeks and 6 weeks after explant preparation and culturing on glyphosate containing medium.

15

Table 4

Glyphosate Level ( $\mu$ M)	Total Number Explants	Number Regenerating 3 Weeks	Number Regenerating 6 Weeks	% Explant Regeneration
<b>Wild Type</b>				
50	10	0	0	0
75	10	0	0	0
100	10	0	0	0
150	10	0	0	0
200	10	0	0	0
<b>38773-6</b>				
50	8	5	8	100
75	18	14	18	100
100	17	12	15	88
150	18	10	16	89
200	16	8	15	86

The above results demonstrate that at all levels of glyphosate examined, shoots regenerated from explants prepared from a tobacco line homoplasmic for pMON38773, while no shoots regenerated from explants prepared from nontransformed control plants. These results suggest that tobacco plants expressing EPSPS in plastids demonstrate tolerance to glyphosate levels of at least 200 $\mu$ M.

Additional transplastomic lines were tested *in vitro* for glyphosate tolerance as described above. The results are shown in Table 5

TABLE 5

Summary of tobacco plastid transformation experiments with various constructs containing EPSPS genes.

5	Construct	Spec/strep (+)	No. of shoots Gly 50 uM(+)
10	pMON38766 (Wild)	1	0
	pMON38766 (T7)	6	0
	pMON38773 (Wild)	9	5 (1)
15	pMON38797 (Wild)	2	0
	pMON38798	6	6
20	pMON38793	8	0
	pMON38796	4	0
	pMON45201	9	3
25	pMON45204	12	*
(No. of shoots positive at 1 mM glyphosate)			

30

These results demonstrate that these transplastomic lines show tolerance to glyphosate. The numbers in parentheses are the number of shoots resistant to selection at 1 mM glyphosate. Thus, as can be seen in table 5, tobacco lines are generated that are tolerant of selection at 1 mM glyphosate.

35

Homoplasmic tobacco plants of the line 38773-6 are sprayed with glyphosate using a track sprayer at concentrations corresponding to 0oz/acre, 16oz/acre, 32oz/acre and 64oz/acre to test for whole plant tolerance. Plant height was measured before and after spraying with glyphosate. The vegetative injury data was collected two weeks after spraying, while the reproductive injury data was collected at plant maturity.

40

Initial results indicate that homoplasmic tobacco lines sprayed are tolerant of glyphosate at the concentration of 16oz/acre as demonstrated in the vegetative tissue injury (Table 6). As can be seen in Table 5 transplastomic lines were generated which

demonstrated a good level of glyphosate tolerance at 32oz/Acre. In subsequent experiments with additional transformed lines, transplastomic lines have shown tolerance to glyphosate at a level of 64oz/Acre.

5 Tolerance is characterized by the continued growth and greening of tissues sprayed with glyphosate. However, as the concentration of glyphosate applied increased, there was a corresponding increase in the level of vegetative injury. In contrast, nontransformed control plants which were highly susceptible to glyphosate concentrations as low as 16 oz/Acre.

10



Table 6

Plant No.	Construct	Roundup rate (oz/A)	Plant height (cm) before spray	Plant height (cm) after spray	Vegetative injury	Fertility rating
1	38773	0	12.2	30.5	0	0
2	38773	0	13.6	34.0	0	0
3	38773	0	8.6	23.8	0	0
4	38773	0	8.6	26.2	0	0
5	38773	0	7.8	28.8	0	0
6	38773	0	12.8	31.5	0	0
7	38773	0	12.2	31.6	0	0
8	38773	0	11.6	35.5	0	0
9	38773	16	9.0	29.0	1	0
10	38773	16	14.4	31.0	0	0
11	38773	16	13.4	32.0	0	0
12	38773	16	13.2	30.0	0	0
13	38773	16	14.2	30.5	0	1
14	38773	16	14.0	33.0	0	0
15	38773	16	13.2	30.2	0	0
16	38773	16	14.9	30.4	0	0
17	38773	32	12.0	26.5	2	4
18	38773	32	11.6	25.4	1	1
19	38773	32	9.4	22.0	1	3
20	38773	32	11.2	23.0	2	4
21	38773	32	13.8	25.8	1	2
22	38773	32	12.4	23.0	1	4
23	38773	32	10.2	19.0	2	4
24	38773	32	13.8	23.2	2	3
26	38773	64	11.8	20.0	2	5
27	38773	64	13.0	22.0	2	5
28	38773	64	12.2	18.0	3	5
29	38773	64	15.8	23.0	2	5
30	38773	64	10.4	17.5	2	5
32	38773	64	15.0	18.5	2	5
33	38773	64	13.8	21.8	2	5
34	38773	64	13.6	19.0	3	5
35	38773	64	10.8	16.0	3	5
36	Wild type	0	21.0	40.6	0	0
37	Wild type	0	16.0	38.0	0	0
38	Wild type	0	15.0	34.6	0	0
39	Wild type	0	17.6	32.2	0	0
40	Wild type	0	15.0	31.6	0	0
41	Wild type	0	14.0	32.0	0	0
42	Wild type	16	10.0	11.8	3	5
43	Wild type	16	8.0	10.0	3	5
44	Wild type	16	8.6	11.0	3	5

45	Wild type	16	8.0	14.0	3	5
46	Wild type	16	9.8	11.0	3	5
47	Wild type	16	10.4	14.0	3	5
48	Wild type	32	10.8	13.2	3	5
49	Wild type	32	9.0	13.0	3	5
50	Wild type	32	8.0	10.2	3	5
51	Wild type	32	11.0	14.0	4	5
52	Wild type	32	9.8	13.0	3	5
53	Wild type	32	8.0	10.8	4	5
54	Wild type	64	7.5	8.6	4	5
55	Wild type	64	11.2	12.5	4	5
56	Wild type	64	10.2	12.8	4	5
57	Wild type	64	11.5	13.0	4	5
58	Wild type	64	13.0	15.0	4	5
59	Wild type	64	9.8	11.2	4	5

Vegetative injuries:

0=normal plant

1=slight chlorosis of new leaves and stunting

2=severe chlorosis of new leaves, malformation of new leaves, and severe stunting

3=dying plant

4=dead plant

Fertility ratings:

0 = Fertile, no delay in maturity, lots of seed

1 = Some abortion, slight delay in seed set, seed

2 = Significant abortion, significant delay in seed set, some seed

3 = Very severe abortion, immature seed pots, a few seed

4 = malformed flowers; if flowered, extreme delay in flowering and no seed produced

5=dead plant

In addition, other transplastomic lines were analyzed for tolerance to spraying with various levels of glyphosate as described above. Specific activity is measured as the amount of exogenously added Phosphoenol pyruvate (PEP) converted to Shikimate-3-phosphate (S3P) per unit protein in the plant extract. Addition of glyphosate tests sensitivity of the EPSPS enzyme to glyphosate. The results are summarized in Table 7.

**Table 7**

LINE	% TOTAL	SPECIFIC	SPECIFIC	Vegetative	Reproductive
------	---------	----------	----------	------------	--------------

	SOLUBLE PROTEIN	ACTIVITY (nmol/min/mg) No gly	ACTIVITY (nmol/min/mg) +1 mM gly	tolerance (oz/acre)	tolerance (oz/acre)
Wild-type		3.4	0	0	0
pMON10154	0.04	19.0	18.4	128	64
pMON45201	-	301.6	221.2	32	32
pMON45204	-	339.9	371.8	128	64
pMON30123	0.001	4.0	0	0	0
pMON30130	0.002	6.2	0	0	0
pMON38773	0.2	16.7	6.7	32	16
pMON38798	0.2	17.2	14.7	32	16
pMON45259	>12.0	-	-	128	64
pMON49218	>12.0	-	-	128	64

These data demonstrate that high levels of glyphosate tolerance can be obtained in transplastomic plants expressing various EPSPS sequences. In particular, lines pMON45204, pMON45259, and pMON49218 provide tolerance to glyphosate applied at levels of at least 128 oz/acre on vegetative tissues, and at least 64 oz/acre on reproductive

5 tissues.

Furthermore, constructs pMON42259 and pMON49218 provide for high level expression of CP4 EPSPS from plant plastids transformed with these constructs. In particular, expression levels of greater than about 12 percent total soluble protein are obtained in constructs employing sequences encoding the first 14 amino acids of GFP

10 fused to the N-terminus of CP4.

### 3F. BT/BXN Analysis

Homoplasmic tobacco plants of the lines 5175 and 5197 are sprayed with Buctril herbicide at a concentration of 4% to test for whole plant tolerance.

Results of the spray test with Buctril demonstrated that all 5197 lines expressing *bxn* were completely resistant when sprayed with a solution containing 4% Buctril herbicide.

Two lines out of six 5175 lines tested were completely resistant to the herbicide when sprayed with a 4% solution containing Buctril.

### 3G. Norflurazon Resistance Analysis

An experiment was set up to determine the efficacy of the *Crt I* trait with respect to resistance to the herbicide Norflurazon. Three 5177 transformed lines, 74-1B-P, 74-2-A, and 74-7-C and three control lines were planted. Plants were grown for seven weeks and then watered with a 3 $\mu$ M Norflurazon solution. Plants negative for the presence of the *crtI* plastid-borne gene were bleached by Norflurazon treatment, positive plants stayed green and continued to grow.

The results show that the three homoplasmic 5177 tobacco lines were resistant to the 3 $\mu$ M Norflurazon solution, while the control plants were all susceptible to the solution (Table 8).

Table 8

Line	Control/Transgenic	Result
Xanthi	Control	Susceptible
2560A Xanthi	Control	Susceptible
75-5D-A	Control	Susceptible
74-1B-P	homoplasmic	Resistant
74-2-A	homoplasmic	Resistant
74-7-C	homoplasmic	Resistant

## 20 Example 4 Analysis of hGH Transgenic Tobacco Plants

### 4A. Southern Analysis

Transformed plants selected for *aadA* marker gene expression are analyzed to determine whether the entire plastid content of the plant has been transformed (homoplastic transformants). Homoplasmic plants are selected using Southern hybridization for further cultivation.

5            Genomic DNA is isolated from transformed tobacco plants, electrophoresed, and transferred to filters as described in Svab *et al.* ((1993), *Proc Natl Acad Sci*, 90:913-917).

             Homoplasmic tobacco plants transformed to express hGH were identified using a probe prepared from a 2.4 kb *EcoRI/EcoRV* fragment from the vector pOVZ2 (similar to pOVZ15 described in Zoubenko, *et al.* 1994, *supra*). The 2.4 kb probe fragment  
10           encompasses part of the targeting sequence.

             The complete disappearance of the 3.27 Kb native tobacco *BamHI* fragment in the lines with a probe covering the region of integration, and the appearance of the expected size band for the inserted DNA fragments in those transformants, 5.6 kb, establishes that the transformed plants are homoplasmic for the intended constructs.

15

#### 4B. Protein Expression Analysis

             Homoplasmic tobacco lines expressing hGH and nuclear tobacco transformants are used to determine the expression of the hGH protein. Western blot analysis was performed on tobacco lines containing constructs pWRG4838, pMON38755 and pMON38794 for  
20           plastid expression and an ELISA assay was used for transgenic tobacco lines containing pWRG4744 and pWRG4747 for nuclear expression of hGH.

             Total protein extractions and western blot procedures were performed as described above, with the exception of the primary antibody was raised against hGH.

Table 9

## Expression Levels of hGH in Tobacco Nuclear Genome and Plastid genome

Construct	Expression	Expression Level % Total Soluble Protein
pWRG4744	nuclear	0.002-0.125%
pWRG4747	nuclear	0.002-0.025%
pWRG4838	plastid	0.2%
pMON38755	plastid	1.0%
pMON38794	plastid	7.0%

Results of the Western analysis (Table 9) demonstrates that hGH expressed in plastids of plant cells accumulates to significantly higher levels than hGH expressed in the nucleus and targeted to either the cytoplasm or plastid of plant cells. Tobacco plants transformed to express hGH in the nucleus accumulated hGH levels of 0.002% (cytoplasmic targeted) to 0.025% (plastid targeted) of total soluble leaf protein, while tobacco plants expressing hGH in the plastid accumulated hGH levels of 0.2% to 7.0% of the total soluble leaf protein as hGH. Furthermore, homoplasmic tobacco plants expressing hGH directed from the *Prm/G10L* regulatory sequences accumulate 35 fold higher levels of hGH than homoplasmic tobacco plants expressing hGH directed from the *PpsbA* promoter sequence.

## 4C. Characterization of hGH Protein Expressed in the Plastid

In order to determine whether the hGH expressed from plastids was properly processed, experiments were performed to determine correct folding and bioactivity.

Two bottom leaves of transplastomic tobacco lines containing pMON38794 were used to extract and purify hGH. Large veins were removed from the excised leaves, and the leaf tissue was cut into small sections (approximately 0.5 cm<sup>2</sup>). The leaf pieces were flash frozen in liquid nitrogen and ground to a fine powder in a chilled mortar and pestle.

Ten grams of frozen, ground leaf tissue was added to ice cold 100 mM Tris base solution (30 ml) and mixed vigorously by vortexing for 5 minutes. The solution was filtered through a single layer of cheese cloth.

From the filtered solution, three separate samples were prepared. The first sample was prepared by centrifuging 4 ml of the filtrate for 1 minute at 16,000 rpm. The centrifugate was aliquoted into 1 ml vials and frozen in dry ice. The remaining filtrate was centrifuged for 10 minutes at 4800 rpm, and several 0.5 ml aliquots were frozen as above for the second sample. To the remaining centrifuged filtrate (approximately 25 ml), 200  $\mu$ l of glacial acetic acid was added to lower the pH from 8.2 to 4.56. The solution was centrifuged at 4800 rpm for 30 minutes, and the supernatant was frozen over dry ice for the third sample.

Total soluble protein (TSP, Table 10) was calculated in these samples by standard protein assay procedures (Maniatis, ), and the percent purity of hGH was calculated based on results from Western blot analysis using known concentrations of starting material.

Table 10

Sample ID	TSP mg/mL	GP2000 mg/L	% Purity
Filtered Extract immediately centrifuged and frozen	6.3	28	0.45 %
Filtered extract centrifuged at 4800 rpm for 10 min and frozen	6.4	28	0.45
pH adjusted and centrifuged extract	0.75	21	2.8%

The pH adjusted and centrifuged extract was purified by Reverse Phase-HPLC (RP-HPLC) for electrospray mass spectrometry and amino-terminal amino acid sequencing. RP-HPLC was performed using a Perkin-Elmer series 200 pump and autosampler and a Vydac C8 (250 by 4.6 mm) RP-HPLC column. 750 microliters of sample was loaded onto the column equilibrated with 20 mM trifluoroacetic acid (TFA) and 50 % acetonitrile. After loading, the column was washed for 2 minutes with 50 %

5 The results of the RP-HPLC analysis are shown in figure 3. Peak I (tallest peak) has the retention time expected for properly folded, native 22 kDa GP2000. This peak was collected and dried down in a Savant Speed-Vac for amino terminal sequencing and electrospray mass spectrometry.

The results of the mass spectrometry of the RP-HPLC peak 1 shows 4 major protein species of different molecular mass. The 21,997 kDa species represents the predicted mass of hGH with the predicted N-terminal Phe removed by over-cleavage of the Ubiquitin protease. The 22,124 kDa species represents the predicted mass of properly processed, correct amino acid sequence of hGH. The 22,507 kDa and 22,664 kDa species are thought to represent an hGH with the N-terminal Phe and hGH which have been modified during plant extraction procedures, respectively. The calculated molecular mass of the proteins suggests that the hGH expressed from the plastid is properly folded (i.e. the correct disulfide bonds are created).

BNSDOCID: <WD 0003022A2 I >



#### 4D. Bioactivity of hGH Expressed in Plant Plastids

Bioactivity of the pH adjusted and centrifuged extract was tested using cells from an Nb2 cell line. These cells proliferate in the presence of growth hormone and other estrogenic type compounds. The assay involves putting various concentrations of growth hormone-containing extract into a 96 well plate. Then a constant amount of cells are added to each well. The plate is incubated for 48hrs and then a reagent called MTS is added. Metabolizing cells take up the MTS and convert it to a blue colored substance. The more cells there are the more blue color in the well. The blue color is measured using a spectrophotometer. The number of cells should be proportional to the concentration of growth hormone in the media. At some high concentration one expects that the cells will become saturated with growth hormone and that the dose response will level off. At very low hGH concentrations essentially no enhanced growth is seen. A sigmoidal shape graph is expected to be produced graphing the cell number (or absorbance) vs hGH concentration graph.

The results of the bioactivity assay (Figure 5) demonstrates that the hGH expressed from a plant plastid has a sigmoidal shape when graphed as absorbance vs hGH concentration.

#### Example 5 Analysis of Aprotinin Transplastomic Tobacco Plants

20

##### 5A. Western Analysis of Aprotinin Expression in Plastids

Homoplasmic tobacco lines expressing are used to determine the expression of the aprotinin protein. Western blot analysis was performed on tobacco lines containing constructs pCGN6146, pCGN6147, pCGN6154 and pCGN6156 for plastid expression of aprotinin.

25

Total protein extractions and western blot procedures were performed as described above, with the exception of the primary antibody was raised against aprotinin.

The results of the Western analysis indicate that aprotinin is expressed from the T7 polymerase promoter when the aprotinin coding sequence is fused with either the PetA or

full length GUS gene. Furthermore, these results indicate that the petA sequence efficiently targets the aprotinin protein to the plant cell thylakoid.

5 All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

10           Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claim.

## CLAIMS

What is claimed is:

- 5
1. A construct comprising the following components in the 5' to 3' direction of transcription:
- 10 a) a promoter functional in a plant plastid;  
b) a DNA sequence which is capable of conferring tolerance in a plant cell to at least one herbicide compound when said DNA sequence is transcribed in plastids of said plant cell; and  
c) a transcription termination region.
- 15 2. The construct according to Claim 1, wherein said construct further comprises (d) a gene encoding a selectable marker for selection of plant cells comprising a plastid expressing said marker and (e) DNA regions of homology to the genome of said plastid, wherein said regions of homology in (e) flank components (a), (b), (c) and (d).
- 20 3. The construct according to Claim 1, wherein said construct further comprises (f) a ribosome binding site joined to said promoter component (a).
- 25 4. The construct according to Claim 3, wherein said ribosome binding site (f) is from a leader sequence selected from the group consisting of sites derived from plastid, bacterial or bacteriophage leader sequences.
5. The construct according to Claim 4, wherein said ribosome binding site is selected from the group consisting of the binding site of the gene 10 leader and the rbcLRBS site.
- 30 6. The construct according to Claim 1 wherein said DNA sequence encodes a gene which confers tolerance to the herbicide glyphosate.
- 35 7. The construct according to Claim 6 wherein said DNA encoding sequence encodes a glyphosate-tolerant 5-enolpyruvylshikimate-3-phosphate synthase.
- 40 8. The construct according to Claim 7 wherein said DNA encoding sequence is selected from the group consisting of the *E. coli* or *Salmonella aroA* gene, the CP4 gene, mutant petunia EPSPS, mutant EPSPS gene of *Psuedomonas* strain LBAA, and the *Bacillus subtilis aroE* gene.
9. The construct according to Claim 6 wherein said DNA encoding sequence encodes a glyphosate-modifying enzyme.
- 45 10. The construct according to Claim 9 wherein said glyphosate-modifying enzyme is selected from the group consisting of the *gox*, *hph*, *glpA* and *glpB*.

11. The construct according to Claim 1 wherein said DNA encoding sequence is the native encoding sequence to said gene.

5 12. The construct according to Claim 1 wherein said DNA encoding sequence is a synthetic encoding sequence to said gene.

13. The construct according to Claim 1 wherein said DNA sequence encodes a sulphonylurea-tolerant AHAS gene.

10 14. The construct according to Claim 1 wherein said DNA sequence encodes a imidazalinone-tolerant AHAS gene.

15 15. The construct according to Claim 14 wherein said DNA encoding sequence is the ALS gene.

16. The construct according to Claim 1 wherein said DNA sequence encodes a phosphinothricin-tolerant gene.

20 17. The construct according to Claim 16 wherein said DNA encoding sequence is the BAR gene.

18. The construct according to Claim 1 wherein said DNA sequence encodes an enzyme of the carotenoid pathway.

25 19. The construct according to Claim 18 wherein said DNA encoding sequence is the crtI gene.

30 20. The construct according to Claim 1 wherein said DNA sequence encodes a bromoxynil-tolerant gene.

21. The construct according to Claim 20 wherein said bromoxynil-tolerant gene is the BXN gene.

35 22. A plant cell plastid containing the construct according to Claim 1.

23. A plant, plant seed, plant cell or progeny thereof containing a plant plastid according to Claim 22.

40 24. A method for producing tolerance of a herbicide in a plant cell, wherein said method comprises transforming plastids of said plant cell with a construct comprising the following as operably joined components in the 5' to 3' direction of transcription:

45 a) a promoter functional in a plant plastid;  
b) a DNA sequence which is capable of conferring in a plant cell tolerance to at least one herbicide compound when said DNA sequence is transcribed in plastids of said plant cell; and

c) a transcription termination region,  
and growing plant cells comprising said transformed plastids under conditions wherein said DNA sequence is transcribed whereby plant cells containing said plant plastids are rendered tolerant to applications of said at least one herbicide compound.

25. The method according to Claim 24, wherein said construct further comprises (d) a gene encoding a selectable marker for selection of plant cells comprising a plastid expressing said marker and (e) DNA regions of homology to the genome of said plastid, wherein said regions of homology in (e) flank components (a), (b), (c) and (d).  
5
26. The method according to Claim 24, wherein said construct further comprises a second DNA sequence of interest.
- 10 27. The method according to Claim 26, wherein said second DNA sequence comprises a gene expressed from an promoter independent from said promoter (a).
28. The method according to Claim 26, wherein said second DNA sequence comprises a gene expressed from said promoter (a) as a polycistronic message with said DNA sequence in (b).  
15
29. The method according to Claim 26, wherein said second DNA sequence comprises a gene other than a gene capable of conferring in a plant cell tolerance to said herbicide compound.  
20
30. The method according to Claim 26, wherein said second DNA sequence comprises a gene capable of conferring in a plant cell tolerance to a second herbicide compound.
- 25 31. The method according to Claim 24, wherein said construct further comprises (f) a ribosome binding site joined to said promoter component (a).
32. The construct according to Claim 31, wherein said ribosome binding site is selected from the group consisting of the binding site of the gene 10 leader and the rbcLRBS site.  
30
33. The method according to Claim 24 wherein said DNA sequence encodes a gene which confers tolerance to the herbicide glyphosate.
- 35 34. The method according to Claim 33 wherein said DNA encoding sequence encodes a glyphosate-tolerant 5-enolpyruvylshikimate-3-phosphate synthase.
35. The method according to Claim 34 wherein said DNA encoding sequence is selected from the group consisting of the E. coli or Salmonella aroA gene, the CP4 gene, mutant petunia EPSPS, mutant EPSPS gene of *Psuedomonas* strain LBAA and the Bacillus subtilis aroE gene.  
40
36. The method according to Claim 33 wherein said DNA encoding sequence encodes a glyphosate-modifying enzyme.  
45
37. The method according to Claim 36 wherein said glyphosate-modifying enzyme is selected from the group consisting of the gox, hph, glpA and glpB.

38. The method according to Claim 24 wherein said DNA encoding sequence is the native encoding sequence to said gene.

5 39. The method according to Claim 24 wherein said DNA encoding sequence is a synthetic encoding sequence to said gene.

40. The method according to Claim 24 wherein said DNA sequence encodes a sulphonylurea-tolerant AHAS gene.

10 41. The method according to Claim 24 wherein said DNA sequence encodes a imidizalinone-tolerant AHAS gene.

15 42. The method according to Claim 41 wherein said DNA encoding sequence is the ALS gene.

43. The method according to Claim 24 wherein said DNA sequence encodes an enzyme of the carotenoid pathway.

20 44. The method according to Claim 43 wherein said DNA encoding sequence is the crtI gene.

45. The method according to Claim 24 wherein said DNA sequence encodes a bromoxynil-tolerant gene.

25 46. The method according to Claim 45 wherein said bromoxynil-tolerant gene is the BXN gene.

47. An herbicide tolerant plant cell produced according to the method of Claim 24.

30 48. A plant, plant seed or plant part comprising a plant cell according to Claim 47.

49. A plant cell according to Claim 47 and comprising greater than about 0.01 % of total soluble protein as a protein expressed from said herbicide-tolerance gene.

35 50. A plant cell according to Claim 47 and comprising greater than about 0.1 % of total soluble protein as a protein expressed from said herbicide-tolerance gene.

40 51. A plant cell according to Claim 47 and comprising greater than about 0.2 % of total soluble protein as a protein expressed from said herbicide-tolerance gene.

52. A plant cell according to Claim 47 and comprising 1 % or more of total soluble protein as a protein expressed from said herbicide-tolerance gene.

45 53. A plant cell according to Claim 47 and comprising 12 % or more of total soluble protein as a protein expressed from said herbicide-tolerance gene.

54. A plant cell according to Claim 47 wherein said herbicide-tolerance gene is a glyphosate-tolerant 5-enolpyruvylshikimate-3-phosphate synthase.

55. A plant, plant seed or plant part comprising a plant cell according to Claim 53.

56. A plant according to method of Claim 55 tolerant of the herbicide glyphosate when said herbicide is applied at a rate of about 16 ounces or greater per acre.

5

57. A plant according to method of Claim 55 tolerant of the herbicide glyphosate when said herbicide is applied at a rate of about 32 ounces or greater per acre.

57. A plant according to method of Claim 55 tolerant of the herbicide glyphosate when said herbicide is applied at a rate of about 64 ounces or greater per acre.

10

58. A plant according to method of Claim 55 tolerant of the herbicide glyphosate when said herbicide is applied at a rate of about 128 ounces or greater per acre.

60. A method wherein herbicide-tolerance produced according to the method of Claim 24 is used to select cells transformed by said construct from non-transformed cells.

15

61. A method according to Claim 60 wherein said herbicide-tolerance gene confers tolerance to the herbicide glyphosate.

20

62. A method according to Claim 61 wherein selection for said herbicide tolerance is made on media containing glyphosate at a concentration from at least about 50  $\mu$ M to about 200  $\mu$ M.

63. A method according to Claim 62 wherein selection for said herbicide tolerance is made on media containing glyphosate at a concentration from about 1 mM.

25

64. A plant cell produced according to the method of Claim 60.

65. The method according to Claim 24, wherein said plant cells are tolerant to applications of herbicide amounts selected from the group consisting of 16 ounces/acre, 32 ounces/acre, 64 ounces/acre, and 128 ounces/acre.

30

66. The method according to Claim 65, wherein said herbicide is glyphosate.


5' -AAT TGT AGA AAT AAT TTT GTT TAA CTT TAA GAA GGA GAT ATA CC-3'  
TTA ACA TCT TTA TTA AAA CAA ATT GAA ATT CTT CCT CTA TAT GG

**FIGURE 1**




Arg-Pro-Asp-Phe-Cys-Leu-Glu-Pro-Pro-Tyr-Thr-Gly-Pro-Cys-  
Lys-Ala-Arg-Ile-Ile-Arg-Tyr-Phe-Tyr-Asn-Ala-Lys-Ala-Gly-  
Leu-Cys-Gln-Thr-Phe-Val-Tyr-Gly-Gly-Cys-Arg-Ala-Lys-Arg-  
Asn-Asn-Phe-Lys-Ser-Ala-Glu-Asp-Cys-Met-Arg-Thr-Cys-Gly-  
Gly-Ala

**FIGURE 2**



RP-HPLC analysis shown above. Peak I (tallest peak) has the retention time expected for properly folded, native 22 kDa GP2000. This peak was collected and dried down in a Savant Speed-Vac for amino terminal sequencing and electrospray mass spectrometry.

**FIGURE 3**



Mass spectrometry of RP-HPLC peak I shown above. Note that there are 4 major protein species, of different molecular mass.

21977 kDa - This is the predicted mass of "des-Phe"-GP2000, ie. GP2000 with the predicted N-terminal Phe removed by over-cleavage of Ubiquitin protease.

\*\*\*\*22124 kDa - This is the predicted mass of properly processed, correct amino acid sequence GP2000

22507 kDa - This is probably des-Phe-GP2000 modified during plant extraction procedures

22664 kDa - This is probably GP2000 modified during plant extraction procedures

Amino terminal sequencing was done by standard Edman degradation, and confirmed the N-terminal sequences discussed above.

**Figure 4**

Summary Graph




FIGURE 5

EcoRI    SacI    KpnI    Prrn  
1   GAATTCGAGC TCGGTACCCA AAGCTCCCC GCCGTCGTT C AATGAGAATG

-35                      -10  
51   GATAAGAGGC TCGTGGGAT GACGTGAGGG GGCAGGGATG GCTATATTTC

PEP                      G10L  
101 TGGGAGCGAA CTCGGGCCA ATTGTAGAA TAATTTTGTT TAACTTTAAG

RBS                      NcoI  
151 AAGGAGATAT ACCCATGG

Figure 6

```

EcoRI      KpnI      Prrn
1  GAATTCGGTA CCCCCGTCGT TCAATGAGAA TGGATAAGAG GCTCGTGGGA

-35      -10      PEP
51  TTGACGTGAG GGGGCAGGGA TGGCTATATT TCTGGGAGCG AACTCCGGGC

NEP
101 GAATACTGAA GCGCTTGGAT ACAAGTTATC CTGGGAAGGA AAGACAATTC

G10L
151 CGGATCCTCT AGAAATAATT TTGTTTAACT TTAAGAAGGA GATATACCC ATG
Met

14aa GFP
GGT AAA GGA GAA GAA CTT TTC ACT GGA GTT GTC CCA
Gly Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro

AGC ATG
Ser Met

```

Figure 7

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/54, 15/55, 15/60, 15/82, 5/10,</b> <b>A01H 5/00, 5/10</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/03022</b> <b>(43) International Publication Date:</b> 20 January 2000 (20.01.00)
<b>(21) International Application Number:</b> PCT/US99/15472 <b>(22) International Filing Date:</b> 10 July 1999 (10.07.99)  <b>(30) Priority Data:</b> 09/113,257 10 July 1998 (10.07.98) US  <b>(71) Applicant:</b> CALGENE LLC [US/US]; 1920 Fifth Street, Davis, CA 95616 (US).  <b>(72) Inventors:</b> HAJDUKIEWICZ, Peter; 15821 Timbervalley Road, Chesterfield, MO 60317 (US). MCBRIDE, Kevin, E.; 1309 Marina Circle, Davis, CA 95616 (US). NEHRA, Narender; 14971 Appalachian Trail, Chesterfield, MO 63017 (US). SCHAAF, David, J.; 2780 Brentwood Place, Davis, CA 95616 (US). STALKER, David, M.; 870 W. Southwood, Woodland, CA 95695 (US). STAUB, Jeffrey, M.; 819 "J" Foxspring Drive, Chesterfield, MO 63017 (US). YE, Guangning; 2325 Crimson View Court, Ellisville, MO 63011 (US).  <b>(74) Agent:</b> MONSANTO/G.D. SEARLE; Patent Dept. Central, P.O. Box 5110, Chicago, IL 60680-5110 (US).	<b>(81) Designated States:</b> BR, CA, JP, MX, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 13 July 2000 (13.07.00)	
<b>(54) Title:</b> EXPRESSION OF HERBICIDE TOLERANCE GENES IN PLANT PLASTIDS  <b>(57) Abstract</b>  Provided are constructs and methods for expressing herbicide tolerance genes in plastids of plant cells. Constructs include the components of a promoter functional in a plant plastid, a DNA sequence which is capable of conferring tolerance in a plant cell to at least one herbicide compound when said DNA sequence is transcribed in plastids of said plant cell and a transcription termination region. Herbicide tolerance is produced by transforming plastids with the constructs of the invention and growing plant cells comprising the transformed plastids under conditions wherein the DNA sequence is transcribed and plant plastids and cells containing the plastids are rendered tolerant to applications of at least one herbicide compound.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/15472

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N15/55 C12N15/60 C12N15/82 C12N5/10  
A01H5/00 A01H5/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 32011 A (CIBA GEIGY AG ;VOLRATH SANDRA L (US); JOHNSON MARIE A (US); POTTER) 4 September 1997 (1997-09-04)  see the whole document; esp. examples 34-38	1-6, 9-27, 29-33, 36-48, 60-64
X	US 5 693 507 A (DANIELL HENRY ET AL) 2 December 1997 (1997-12-02)  see the whole document; esp. example 4  -/-	1-27, 29-48, 54,60-64

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

2 May 2000

Date of mailing of the international search report

10. 05. 2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3018

Authorized officer

Kania, T

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/15472

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DANIELL ET AL: "Containment of herbicide resistance through genetic engineering of the chloroplast genome" NATURE BIOTECHNOLOGY, US, NATURE PUBLISHING, vol. 16, April 1998 (1998-04), page 345-348 XP002090409 ISSN: 1087-0156 the whole document	1-27, 29-48, 54, 60-64
X	DATTA ET AL: "Transformation of the tobacco chloroplast genome with the aroA gene to confer glyphosate tolerance" PLANT PHYSIOLOGY, US, AMERICAN SOCIETY OF PLANT PHYSIOLOGISTS, ROCKVILLE, MD, vol. 111, no. 2, 1996, page 168 XP002090406 ISSN: 0032-0889 abstract	1-8, 11, 12, 22-25, 31-35, 38, 39, 47, 48, 54, 60-64
A	MCBRIDE ET AL: "Amplification of a chimeric Bacillus gene in chloroplasts leads to an extraordinary level of an insecticidal protein in tobacco" BIO/TECHNOLOGY, US, NATURE PUBLISHING CO. NEW YORK, vol. 13, 1995, page 362-365 XP002090404 ISSN: 0733-222X the whole document	1-8, 11, 12, 22-25, 31-35, 38, 39, 47-66
A	WO 98 11235 A (CIBA GEIGY AG ; HEIFETZ PETER (US); LEBEL EDOUARD (US); UKNES SCOTT) 19 March 1998 (1998-03-19)  see example C	1-8, 11, 12, 22-25, 31-35, 38, 39, 47-66
A	WO 95 24493 A (CALGENE INC ; MCBRIDE KEVIN E (US); STALKER DAVID M (US)) 14 September 1995 (1995-09-14)  the whole document	1-8, 11, 12, 22-25, 31-35, 38, 39, 47-66
A	WO 95 16783 A (CALGENE INC ; MCBRIDE KEVIN E (US); STALKER DAVID M (US)) 22 June 1995 (1995-06-22) cited in the application  the whole document	1-8, 11, 12, 22-25, 31-35, 38, 39, 47-66

-/--

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/15472

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 293 358 A (MONSANTO CO) 30 November 1988 (1988-11-30)  the whole document	1-8,11, 12, 22-25, 31-35, 38,39, 47-66
A	WO 92 04449 A (MONSANTO CO) 19 March 1992 (1992-03-19) cited in the application  the whole document	1-8,11, 12, 22-25, 31-35, 38,39, 47-66
A	WO 97 04103 A (LEBRUN MICHEL ; SAILLAND ALAIN (FR); FREYSSINET GEORGES (FR); RHONE) 6 February 1997 (1997-02-06)  the whole document	1-8,11, 12, 22-25, 31-35, 38,39, 47-66
A	US 5 589 612 A (JILKA JOSEPH M ET AL) 31 December 1996 (1996-12-31) see esp. p.10 l.9-26; p.14 l.48-60	60,61,64
A	ZHOU H ET AL: "GLYPHOSATE-TOLERANT CP4 AND GOX GENES AS A SELECTABLE MARKER IN WHEAT TRANSFORMATION" PLANT CELL REPORTS,DE,SPRINGER VERLAG, vol. 15, no. 3/04, 1 December 1995 (1995-12-01), pages 159-163, XP000578832 ISSN: 0721-7714 the whole document	61-64
A	US 5 633 437 A (BERNASCONI PAUL ET AL) 27 May 1997 (1997-05-27)  see the whole document; esp. p.5 l.14-34  -/-	1-5, 11-15, 22-27, 29-32, 38-42, 47-53, 55,65

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/15472

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MISAWA N ET AL: "FUNCTIONAL EXPRESSION OF THE ERWINIA UREDOVORA CAROTENOID BIOSYNTHESIS GENE CRTL IN TRANSGENIC PLANTS SHOWING AN INCREASE OF BETA-CAROTENE BIOSYNTHESIS ACTIVITY AND RESISTANCE TO THE BLEACHING HERBICIDE NORFLURAZON" PLANT JOURNAL, GB, BLACKWELL SCIENTIFIC PUBLICATIONS, OXFORD, vol. 4, no. 5, 1 January 1993 (1993-01-01), pages 833-840, XP002017202 ISSN: 0960-7412 the whole document	1-5, 11, 12, 18, 19, 22-25, 31, 32, 38, 39, 43, 44, 47-53, 55, 65
A	EP 0 337 899 A (RHONE POULENC AGROCHIMIE) 18 October 1989 (1989-10-18)  the whole document	1-5, 11, 12, 20, 22-25, 31, 32, 38, 39, 45, 47-53, 55, 65
A	WO 98 20144 A (HAWKES TIMOTHY ROBERT ;KNIGHT MARY ELIZABETH (GB); THOMPSON PAUL A) 14 May 1998 (1998-05-14)  see the whole document; esp. examples 4-9	1-6, 9-27, 29-33, 36-53, 55-58, 65, 66
P, X	WO 99 05265 A (BAILEY ANA MARIA ;INVEST Y DE ESTUDIOS AVANZADOS (MX); BLOWERS ALA) 4 February 1999 (1999-02-04)  the whole document	1-6, 9-12, 16, 17, 22-33, 36-39, 47, 48, 56, 65, 66

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 99/15472

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
  
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99/15472

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-6,11,12,22-25,31-33,38,39,47-53,  
55-66 partially; 7,8,34,35,54 completely

Plant plastid expression constructs comprising: a promoter functional in a plant plastid, a DNA sequence, native or synthetic, capable of conferring tolerance in the plant cell to at least one herbicide compound, and a transcription termination region. Said constructs further comprising a selectable marker as specified, DNA regions of homology to the plastid genome flanking the other component, a ribosome binding site, preferably the binding site of the gene 10 leader or rbcLRBS. Said DNA sequence encoding a gene which confers tolerance to glyphosate, preferably an EPSPS encoding gene out of a selection as claimed. Plant cell plastids, plant cells, plants, and seeds comprising said constructs.

A method for producing tolerance of a herbicide in a plant cell by transforming plant plastids with said constructs and growing plant cells under conditions wherein said DNA sequence is transcribed and the plant cells are rendered tolerant to herbicide applications. Said method employing the said constructs as defined above.

Plants and plant cells derived therefrom comprising an amount of said protein as claimed, and tolerant to herbicide concentrations as claimed.

A method wherein herbicide-tolerance produced according to said method is used to select transformed cells from non-transformed cells under selection conditions as defined, and plant cell produced thereby.

2. Claims: 1-6,11,12,22-25,31-33,38,39,47-53,  
55-66 partially; 9,10,36,37 completely

idem, the DNA sequence encoding a glyphosate-modifying enzyme out of a selection as claimed

3. Claims: 1-5,11,12,22-25,31,32,38,39,47-53,55,60,64,  
65 partially; 13-15,40-42 completely

idem, said DNA sequence encoding a sulphonylurea-tolerant AHAS gene, a imidazalinone-tolerant AHAS gene, preferably a ALS gene

4. Claims: 1-5,11,12,22,23 partially; 16,17 completely

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99/15472

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Plant plastid expression constructs comprising: a promoter functional in a plant plastid, a DNA sequence, native or synthetic, capable of conferring tolerance in the plant cell to at least one herbicide compound, and a transcription termination region. Said constructs further comprising a selectable marker as specified, DNA regions of homology to the plastid genome flanking the other component, a ribosome binding site, preferably the binding site of the gene 10 leader or rbcLRBS. Said DNA sequence encoding a phosphinotricin-tolerant gene, preferably the BAR gene. Plant cell plastids, plant cells, plants, and seeds comprising said constructs.

5. Claims: 1-5, 11, 12, 22-25, 31, 32, 38, 39, 47-53, 55, 60, 64, 65 partially; 18, 19, 43, 44 completely

as subject 3, the DNA sequence encoding an enzyme of the carotenoid pathway, preferably the crtI gene

6. Claims: 1-5, 11, 12, 22-25, 31, 32, 38, 39, 47-53, 55, 60, 64, 65 partially; 20, 21, 45, 46 completely

idem, the DNA sequence encoding a bromoxynil-tolerant gene, preferably the BXN gene

7. Claims: 24 partially; 26-30 completely

A method for producing tolerance of a herbicide in a plant cell by transforming plant plastids with a construct as defined and growing plant cells under conditions wherein said DNA sequence is transcribed and the plant cells are rendered tolerant to herbicide applications.

The method wherein said construct further comprises a second DNA sequence of interest. Said DNA sequence optionally being expressed independently from said promoter, or together with the first DNA sequence from the same promoter as a polycistronic message. Said DNA sequence comprising a gene other than a gene capable of conferring in a plant cell tolerance to said herbicide compound, or capable of conferring tolerance to a second herbicide compound.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/15472

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9732011 A	04-09-1997	AU 1984697 A	16-09-1997
		AU 2065497 A	16-09-1997
		BR 9707769 A	27-07-1999
		BR 9707783 A	27-07-1999
		CA 2247074 A	04-09-1997
		CA 2247797 A	04-09-1997
		CN 1212724 A	31-03-1999
		CN 1212725 A	31-03-1999
		CZ 9802726 A	16-12-1998
		CZ 9802727 A	11-11-1998
		EP 0883682 A	16-12-1998
		EP 0885305 A	23-12-1998
		HU 9900623 A	28-06-1999
		HU 9901044 A	28-07-1999
		PL 328617 A	01-02-1999
		PL 328651 A	15-02-1999
		WO 9732028 A	04-09-1997
		US 5939602 A	17-08-1999
US 5693507 A	02-12-1997	US 5932479 A	03-08-1999
WO 9811235 A	19-03-1998	AU 4414697 A	02-04-1998
		CN 1230224 A	29-09-1999
		EP 0925362 A	30-06-1999
		PL 331767 A	02-08-1999
WO 9524493 A	14-09-1995	US 5545817 A	13-08-1996
		CA 2185339 A	14-09-1995
		EP 0749491 A	27-12-1996
		JP 9510100 T	14-10-1997
		US 5866421 A	02-02-1999
WO 9516783 A	22-06-1995	US 5576198 A	19-11-1996
		CA 2178645 A	22-06-1995
		EP 0734446 A	02-10-1996
		JP 9510342 T	21-10-1997
EP 0293358 A	30-11-1988	US 4971908 A	20-11-1990
		AT 108203 T	15-07-1994
		AU 1660188 A	01-12-1988
		CN 1032030 A	29-03-1989
		DE 3850527 D	11-08-1994
		DE 3850527 T	22-12-1994
		DK 285688 A	05-01-1989
		ES 2058338 T	01-11-1994
		JP 1039984 A	10-02-1989
		NZ 224782 A	26-02-1991
		US 5145783 A	08-09-1992
		US 5312910 A	17-05-1994
		ZA 8803735 A	26-04-1989
WO 9204449 A	19-03-1992	AT 139566 T	15-07-1996
		AU 655945 B	19-01-1995
		AU 8657891 A	30-03-1992
		CA 2088661 A	01-03-1992
		DE 69120419 D	25-07-1996
		DE 69120419 T	09-01-1997
		DK 546090 T	29-07-1996



# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No  
PCT/US 99/15472

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9204449 A		EP 0546090 A	16-06-1993
		ES 2089232 T	01-10-1996
		GR 3021058 T	31-12-1996
		JP 6500472 T	20-01-1994
		US 5633435 A	27-05-1997
		US 5627061 A	06-05-1997
		US 5804425 A	08-09-1998
WO 9704103 A	06-02-1997	FR 2736926 A	24-01-1997
		AU 6619196 A	18-02-1997
		BG 102247 A	30-04-1999
		BR 9609792 A	16-03-1999
		CA 2223875 A	06-02-1997
		CN 1196088 A	14-10-1998
		CZ 9800174 A	17-06-1998
		EP 0837944 A	29-04-1998
		HU 9900463 A	28-05-1999
		JP 11510043 T	07-09-1999
		PL 324572 A	08-06-1998
		SK 6498 A	04-11-1998
US 5589612 A	31-12-1996	NONE	
US 5633437 A	27-05-1997	NONE	
EP 0337899 A	18-10-1989	FR 2629098 A	29-09-1989
		AT 188250 T	15-01-2000
		AU 629539 B	08-10-1992
		AU 3159189 A	28-09-1989
		BR 8901499 A	14-11-1989
		CA 1314505 A	16-03-1993
		CN 1036038 A,B	04-10-1989
		DD 294040 A	19-09-1991
		DE 68929125 D	03-02-2000
		DK 144689 A	24-09-1989
		EG 20026 A	30-10-1997
		EP 0894865 A	03-02-1999
		HU 52158 A,B	28-06-1990
		IL 89614 A	19-01-1996
		JP 2009374 A	12-01-1990
		KR 145703 B	01-08-1998
		NZ 228458 A	25-06-1991
		PL 278448 A	27-12-1989
		RO 107001 A	30-08-1993
		TR 23682 A	06-06-1990
		US 5559024 A	24-09-1996
		US 5773698 A	30-06-1998
		ZA 8902146 A	29-11-1989
WO 9820144 A	14-05-1998	AU 4789597 A	29-05-1998
		BG 103462 A	31-01-2000
		BR 9712695 A	19-10-1999
		CN 1236394 A	24-11-1999
		CZ 9901616 A	11-08-1999
		EP 0946737 A	06-10-1999
		PL 333131 A	22-11-1999
WO 9905265 A	04-02-1999	AU 8583798 A	16-02-1999



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 January 2000 (20.01.2000)

PCT

(10) International Publication Number  
WO 00/03022 A3

- (51) International Patent Classification<sup>7</sup>: C12N 15/54, 15/55, 15/60, 15/82, 5/10, A01H 5/00, 5/10  
Chesterfield, MO 63017 (US). YE, Guangning; 2325  
Crimson View Court, Ellisville, MO 63011 (US).
- (21) International Application Number: PCT/US99/15472 (74) Agent: SCHWEDLER, Carl, J.; Legal/Patent Dept., Cal-  
gene LLC, 1920 Fifth Street, Davis, CA 95616 (US).
- (22) International Filing Date: 10 July 1999 (10.07.1999) (81) Designated States (national): BR, CA, JP, MX.
- (25) Filing Language: English (84) Designated States (regional): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE).
- (26) Publication Language: English
- (30) Priority Data: 09/113,257 10 July 1998 (10.07.1998) US  
Published:  
— With international search report.
- (71) Applicant: CALGENE LLC [US/US]; 1920 Fifth Street,  
Davis, CA 95616 (US). (88) Date of publication of the international search report:  
13 July 2000
- (72) Inventors: HAJDUKIEWICZ, Peter; 15821 Timber-  
valley Road, Chesterfield, MO 60317 (US). MCBRIDE,  
Kevin, E.; 1309 Marina Circle, Davis, CA 95616 (US).  
NEHRA, Narender; 14971 Appalachian Trail, Chester-  
field, MO 63017 (US). SCHAAF, David, J.; 2780  
Brentwood Place, Davis, CA 95616 (US). STALKER,  
David, M.; 870 W. Southwood, Woodland, CA 95695  
(US). STAUB, Jeffrey, M.; 819 "J" Foxspring Drive,  
(48) Date of publication of this corrected version:  
5 July 2001
- (15) Information about Correction:  
see PCT Gazette No. 27/2001 of 5 July 2001, Section II
- For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

WO 00/03022 A3

(54) Title: EXPRESSION OF HERBICIDE TOLERANCE GENES IN PLANT PLASTIDS

(57) Abstract: Provided are constructs and methods for expressing herbicide tolerance genes in plastids of plant cells. Constructs include the components of a promoter functional in a plant plastid, a DNA sequence which is capable of conferring tolerance in a plant cell to at least one herbicide compound when said DNA sequence is transcribed in plastids of said plant cell and a transcription termination region. Herbicide tolerance is produced by transforming plastids with the constructs of the invention and growing plant cells comprising the transformed plastids under conditions wherein the DNA sequence is transcribed and plant plastids and cells containing the plastids are rendered tolerant to applications of at least one herbicide compound.

## EXPRESSION OF HERBICIDE TOLERANCE GENES IN PLANT PLASTIDS

5

### INTRODUCTION

#### Technical Field

This invention relates to the application of genetic engineering techniques to  
10 plants. Specifically, the invention relates to compositions and methods for enhancing  
expression of proteins in plant plastids.

#### Background

The plastids of higher plants are an attractive target for genetic engineering. Plant  
15 plastids (chloroplasts, amyloplasts, elaioplasts, etioplasts, chromoplasts, etc.) are the major  
biosynthetic centers that, in addition to photosynthesis, are responsible for production of  
industrially important compounds such as amino acids, complex carbohydrates, fatty acids,  
and pigments. Plastids are derived from a common precursor known as a proplastid and  
thus the plastids present in a given plant species all have the same genetic content. Plant  
20 cells contain 500-10,000 copies of a small 120-160 kilobase circular genome, each  
molecule of which has a large (approximately 25kb) inverted repeat. Thus, it is possible to  
engineer plant cells to contain up to 20,000 copies of a particular gene of interest which  
potentially can result in very high levels of foreign gene expression. In addition, plastids  
of most plants are maternally inherited. Consequently, unlike heterologous genes  
25 expressed in the nucleus, heterologous genes expressed in plastids are not pollen  
disseminated, therefore, a trait introduced into a plant plastid will not be transmitted to  
wild-type relatives.

There remains a need for improved regulatory elements for expression of genes in a  
plant plastid. To date, the expression signals used routinely for plastid transgene

expression derive from endogenous plastid genes. The plastid expression signals are typically derived from promoter regions of highly expressed plastid genes such as the promoter regions from the 16S ribosomal RNA operon (*Prm*), *psbA* gene (*PpsbA*) or the *rbcl* gene (*Prbcl*). The *psbA* and *rbcl* genes are highly transcribed, but their translation is  
5 controlled by tissue-specific and light-regulated factors which limits their usefulness. In the case of *Prm*, a synthetic ribosome binding site (RBS) patterned after the plastid *rbcl* gene leader has been typically used to direct translation. However, this *Prm*/RBS is translated inefficiently due to poor ribosome binding.

A totally heterologous expression system has been used to express plastid genes  
10 (USPN 5,576,198, the entirety of which is incorporated herein by reference). This system is a two component system. The first component is a plastid transgene driven by a T7 bacteriophage gene 10 promoter/leader sequence. The second component is a nuclear gene encoding the T7 Polymerase that is targeted to the plastid compartment. The limitation of this system is the need to create nuclear transformed lines that express the T7 Polymerase  
15 in preferred ways.

Plastids of higher plants present an attractive target for genetic engineering. As mentioned above, plastids of higher plants are maternally inherited. This offers an advantage for genetic engineering of plants for tolerance or resistance to natural or chemical conditions. such as herbicide tolerance, as these traits will not be transmitted to  
20 wild-type relatives. In addition, the high level of foreign gene expression is attractive for engineered traits such as the production of pharmaceutically important proteins.

Expression of nucleic acid sequences encoding for enzymes providing for herbicide tolerance as well as pharmaceutical proteins from plant plastid genome offers an attractive alternative to expression from the plant nuclear genome.

25

## SUMMARY OF THE INVENTION

The present invention provides nucleic acid sequences useful in enhancing expression of a wide variety of genes, both eukaryotic and prokaryotic, in plant plastids.

Furthermore, plastid expression constructs are provided which are useful for genetic engineering of plant cells and which provide for enhanced expression of the EPSP synthase proteins or the hGH protein in plant cell plastids. The transformed plastids should be metabolically active plastids, and are preferably maintained at a high copy number in the plant tissue of interest, most preferably the chloroplasts found in green plant tissues, such as leaves or cotyledons.

The plastid expression constructs for use in this invention generally include a plastid promoter region capable of providing for enhanced expression of a DNA sequence, a DNA sequence encoding an EPSPS protein or hGH, and a transcription termination region capable of terminating transcription in a plant plastid.

The plastid promoter region of the present invention is preferably linked to a ribosome binding site which provides for enhanced translation of mRNA transcripts in a plant plastid.

The plastid expression construct of this invention is preferably linked to a construct having a DNA sequence encoding a selectable marker which can be expressed in a plant plastid. Expression of the selectable marker allows the identification of plant cells comprising a plastid expressing the marker.

In a preferred embodiment, vectors for transfer of the construct into a plant cell include means for inserting the expression and selection constructs into the plastid genome. This preferably comprises regions of homology to the target plastid genome which flank the constructs.

The constructs of the present invention preferably comprises a promoter sequence linked to a ribosome binding site capable of enhancing the translation of mRNA transcripts in the plant plastid. The ribosome binding site is preferably derived from the T7 bacteriophage gene 10 leader sequence.

Of particular interest in the present invention is the high level of expression of nucleic acid sequences in plant plastids. Of particular interest is the high level expression of nucleic acid sequences encoding for enzymes involved in herbicide tolerance and encoding for pharmaceutical proteins.

The constructs of the present invention preferably comprises a DNA sequence encoding for a 5-Enolpyruvylshikimate-3-phosphate synthase (USPN 5,633,435, the entirety of which is incorporated herein by reference), nitrilase, phytoene desaturase, aprotonin or a DNA sequence encoding Human Growth Hormone (USPN 5,424,199, the  
5 entirety of which is incorporated herein by reference).

Plant cell plastids containing the constructs are also contemplated in the invention, as are plants, plant seeds, plant cells or progeny thereof containing plastids comprising the construct.

The present invention also includes methods for enhanced expression of DNA  
10 sequences in plant plastids.

The invention also includes a method for the enhanced expression of an enzyme conferring herbicide tolerance in a plant cell, by expressing the *Agrobacterium tumefaciens* sp stain CP4 EPSPS in plastids of the plant cell.

In addition, the invention also includes a method for the enhanced expression of an  
15 enzyme encoding hGH in plastids of the plant cell.

Thus, the present invention relates to a chimeric gene containing a herbicide tolerance coding sequence or the coding sequence of a pharmaceutical protein, a plant plastid expression vector containing a promoter operably linked to a T7 Bacteriophage Polymerase gene 10 ribosome binding site capable of enhanced expression in a plant  
20 plastid operably linked to a herbicide tolerance or pharmaceutical coding gene, a plant transformation vector having inserted therein a herbicide tolerance or pharmaceutical coding gene expressed from a plastid promoter linked to a T7 Bacteriophage Polymerase gene 10 ribosome binding site , plant cells transformed using such vectors and plants regenerated therefrom which exhibit a substantial degree of expression of nucleic acid  
25 sequences and proteins and methods for producing such plants and such plants.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide sequence of the G10L ribosome binding site.

Figure 2 provides an amino acid sequence encoding for aprotinin.

5        Figure 3 provides the results of RP-HPLC analysis for characterization of hGH protein expressed in the plastid. Peak I (tallest peak) indicates the expected retention time for properly folded, native 22 kDa GP2000.

Figure 4 provides an electrospray ionization mass spectrometry (MS) analysis using a Micromass Q-Tof electrospray time-of-flight mass spectrometer. In particular, a  
10        series of ions corresponding to the specie(s) present in the sample with varying numbers of protons attached is provided. The axes of the spectrum are intensity versus mass-to-charge ratio of the specie(s) present.

Figure 5 provides a graphic representation of the bioactivity of hGH expressed from a plant plastid. The samples represented on the graph are bovine prolactin (bPL),  
15        hGH expressed from *E. coli* (Ala-hGH), and a null transgenic spiked with bovine prolactin (SPFF Null Spike) as positive controls, a null transgenic (SPFF Null) as a negative control, and transgenic samples from a sepharose column (SPFF Sample, SPFF Sample) and a transgenic sample eluted from the sepharose column at pH3.5 (SPFF pH3.5 Eln).

Figure 6 provides the nucleic acid sequence for the Prn/G10L promoter/RBS  
20        hybrid. The Prn promoter contains the consensus plastid -35 and -10 promoter elements (underlined) and the transcription start sites (GC in bold) for the Plastid-Encoded RNA Polymerase (PEP). The gene 10 leader (G10L) contains a perfect plastid ribosome binding site (RBS, nucleotides in bold).

Figure 7 provides the nucleic acid sequence for the Prn/NEP/G10L::14aaGFP  
25        fusion. The NEP promoter region is underlined (A in bold is transcription start site). The NEP promoter region used extends beyond the consensus sequence both upstream and downstream of the promoter. The initial ATG, the initiator methionine is not counted in the 14 amino acids of GFP.

30

## DETAILED DESCRIPTION OF THE INVENTION

In accordance with the subject invention, plastid expression constructs are provided which generally comprise a promoter functional in a plant plastid, a ribosome binding site



derived from the T7 Bacteriophage Polymerase gene 10 leader, a DNA sequence encoding for a gene of interest, and a transcription termination region capable of terminating transcription in a plant plastid. These elements are provided as operably joined components in the 5' to 3' direction of transcription.

- 5           Furthermore, the constructs of the present invention may also include a nucleic acid sequence encoding a peptide capable of targeting said DNA sequence encoding a protein to the thylakoid lumen within the chloroplast.

Of particular interest in the present invention is the use of the plastid expression constructs to direct the high level transcription and translation (expression) of nucleic acid  
10 sequences. Such plastid expression constructs find use in directing the high level expression of DNA sequences encoding for enzymes involved in herbicide tolerance or encoding for the production of pharmaceutical proteins.

Of more particular interest in the present invention is the use of the plastid expression constructs to direct the high level translation of transcribed messenger RNA.

- 15           DNA sequence and biochemical data reveal a similarity of the plastid organelle's transcriptional and translational machineries and initiation signals to those found in prokaryotic systems. In fact, plastid derived promoter sequences have been reported to direct expression of reporter genes in prokaryotic cells. In addition, plastid genes are often organized into polycistronic operons as they are in prokaryotes.

- 20           Despite the apparent similarities between plastids and prokaryotes, there exist fundamental differences in the methods used to control gene expression in plastids and prokaryotes. As opposed to the transcriptional control mechanisms typically observed in prokaryotes, plastid gene expression is controlled predominantly at the level of translation and mRNA stability by trans-acting nuclear encoded proteins.

- 25           Translation is a multi-stage process which first involves the binding of messenger RNA (mRNA) to ribosomes. Beginning at the translation start codon, the mRNA codons are read sequentially as the ribosomes move along the mRNA molecule. The specified amino acids are then sequentially added to the growing polypeptide chain to yield the protein or polypeptide encoded in the mRNA.

As mentioned, the first step in the translation process is the binding of the mRNA molecule to the ribosome. The nature of this interaction (i.e. binding) has been only partially elucidated. Analysis of RNase-resistant oligonucleotides isolated from bacterial translation initiation complexes indicate that a RNA fragment approximately 30 to 40  
5 nucleotides in length comprises the initial ribosome binding site (RBS). Thus, a RBS is hereinafter understood to comprise a sequence of mRNA surrounding the translation start codon which is responsible for the binding of the ribosome and for initiation of translation.

Recently, ribosome binding sites have been identified capable of directing translation in a prokaryotes. For example, a ribosome binding site derived from the T7  
10 bacteriophage gene 10 leader, G10L (USPN 5,232,840, the entirety of which is incorporated herein by reference), has been identified which enhances expression of nucleic acid sequences in prokaryotes.

Herbicides such as N-phosphonomethylglycine, halogenated hydroxybenzonitriles, and norflurazon have been the subject of a large amount of investigation.

15 N-phosphonomethylglycine, commonly referred to as glyphosate, inhibits the shikimic acid pathway which leads to the biosynthesis of aromatic compounds including amino acids, plant hormones and vitamins. Specifically, glyphosate curbs the conversion of phosphoenolpyruvic acid (PEP) and 3-phosphoshikimic acid to 5-enolpyruvyl-3-phosphoshikimic acid by inhibiting the enzyme 5-enolpyruvylshikimate-3-phosphate  
20 synthase (hereinafter referred to as EPSP synthase or EPSPS).

Glyphosate tolerant plants have been produced by transformation of various EPSP synthase genes into the nuclear genome of a plant. A gene for EPSP synthase has been cloned from *Agrobacterium tumefaciens* sp strain CP4 (USPN 5,633,435) and confers a high level of glyphosate tolerance in plants. Furthermore, high levels of glyphosate  
25 tolerance has been achieved in a number of crop plants by fusing EPSPS to a chloroplast transit peptide (CTP) for targeted expression in plastids. In addition, variants of the wild-type EPSPS enzyme have been isolated which are glyphosate tolerant as a result of alterations in the EPSPS amino acid coding sequence (Kishore and Shah, *Ann. Rev. Biochem.* (1988) 57:627-663; Shulze et al., *Arch. Microbiol.* (1984) 137:121-123; Kishore

5 et al., *FEBS Lett.* (1984) 173: 238-241; Shulze et al., *Arch. Microbiol.* (1984) 137:121-123; Kishore et al., *Fed. Proc.* (1986) 45:1506; Sost and Amrhein, *Arch. Biochem. Biophys.* (1990) 282: 433-436).

10 hydroxybenzonitriles, and norflurazon using nucleic acid sequences expressed in the  
nucleus.

inhibiting electron transfer (Van Rensen (1982) *Physiol. Plant* 54:515-520, and Sanders  
15 and Pallett (1986) *Pestic. Biochem. Physiol.* 26:116-122). Herbicides such as norflurazon  
inhibit the production of carotenoids.

20 bacteria such as *Klebsiella pneumoniae* and used to construct vectors to direct the  
expression of the DNA sequence in plant cell nucleus (USPN 4,810,648, the entirety of  
which is incorporated herein by reference).

25 in plant cell nuclei. For example, by expressing a phytoene desaturase from *Erwinia*  
*uredovora* provides tolerance to norflurazon.

While plants transformed to express nucleic acid sequences encoding for such enzymes from the nuclear genome have found utility in engineering herbicide tolerant

plants, it would be increasingly beneficial to obtain herbicide tolerant plants via plastidial expression.

In the examples provided herein, DNA sequences encoding for enzymes involved in herbicide tolerance are used in constructs to direct the expression of the sequences from the plant plastid. DNA sequences encoding for 5-enolpyruvylshikimate-3-phosphate  
5    synthase (EPSPS), bromoxynil nitrilase (Bxn), phytoene desaturase (crtI (Misawa et al, (1993) *Plant Journal* 4:833-840, and (1994) *Plant Jour* 6:481-489), and acetohydroxyacid synthase (AHAS (Sathasiivan et al. (1990) *Nucl. Acids Res.* 18:2188-2193)) are used in the expression constructs of the present invention to direct the expression of said herbicide  
10    tolerance nucleotide sequences from the plant plastid.

Transplastomic tobacco plants are identified which are homoplasmic for the DNA sequences of interest encoding said herbicide tolerance genes. Homoplasmic plants demonstrate a high level of protein expression from the plastid. Furthermore, homoplasmic plants demonstrate a high level of tolerance for the respective herbicide. For  
15    example, as described in more detail in the example below, plants transformed to express EPSPS from the plastid demonstrate a high level of tolerance for the herbicide glyphosate. In addition, homoplasmic tobacco lines expressing nitrilase or phytoene desaturase demonstrate high levels of tolerance for the herbicides bromoxynil and norflurazon, respectively.

20    An artisan skilled in the art to which the present invention pertains will recognize that additional sequences may be employed to in the plastid expression constructs of the instant invention to produce herbicide tolerant plants. Other nucleic acid sequence which may find use in the plastid expression constructs herbicide tolerant plants include the *bar* gene for tolerance to glufosinate (DeBlock, et al. (1987) *EMBO J.* 6:2513-2519).

25    Furthermore, additional glyphosate tolerance genes may be employed in the constructs of the present invention. Additional glyphosate tolerant EPSPS genes are described in U.S. Patent Number 5,627,061, Padgett et al. (1996) *Herbicide Resistant Crops*, Lewis Publishers, 53-85, and in Penaloza-Vazquez, et al. (1995) *Plant Cell Reports* 14:482-487, the entireties of which are incorporated herein by reference.

It should be noted that the herbicide tolerance constructs of the present invention may also include sequences encoding genes involved in other stress tolerance genes, for example insect or disease resistance/tolerance genes. As described in more detail in the examples that follow, plastid expression constructs are used to regenerate plants which are  
5 resistant to the herbicide Buctril, which also expresses the *Bacillus thuringensis cry1Ac* protein.

In addition, the plastid expression constructs also find use in directing the production of human biological proteins (pharmaceutical proteins) from the plant plastid. Nucleic acid sequences encoding for the Human Growth Hormone (hGH) are employed in  
10 the plastid expression constructs of the present invention. Furthermore, transplastomic tobacco plants containing such constructs demonstrate a high level of expression of hGH. In addition, the hGH protein expressed from the plant plastid exhibits characteristics of proper processing as well as proper protein folding.

Traditional methods of pharmaceutical protein production generally employ  
15 prokaryotic or single cell eukaryotic organisms for expression and large scale production systems. For example, production of the human biologic, Human Growth Hormone (USPN 5,424,199), has been achieved in *Bacillus* and *E. coli* cells.

Another example is the production of aprotinin. Traditional methods for the production of aprotinin have employed the expression of aprotinin in bacteria, or more  
20 usually, the extraction of the protein from bovine organs or tissues. Thus, there is a need in the art for an alternative approach for the large scale production of such human biologics.

Human Growth Hormone (hGH) participates in much of the regulation of normal human growth and development. This 22,000 dalton pituitary hormone exhibits a  
25 multitude of biological effects including linear growth (somatogenesis), lactation, activation of macrophages, insulin-like and diabetogenic effects among others (Chawla, *Ann. Rev. Med.* (1983) 34:519; Edwards, *et al.*, *Science* (1988) 239:769; Thorner *et al.*, *J. Clin. Invest.* (1988) 81:745). Growth deficiency in children leads to dwarfism, which has been successfully treated for more than a decade by exogenous administration of hGH.

hGH is a member of a family of homologous hormones that include placental lactogens, prolactins, and other genetic and species variants or growth hormone (Nicoll, *et al.*, *Endocrine Reviews*(1986) 7:169). hGH is unusual among these in that it exhibits broad species specificity and binds to either the cloned somatogenic (Leung, *et al.*, *Nature* (1987) 33:537) or prolactin receptor (Boutin, *et al.*, *Cell* (1988) 53:69).

Aprotinin (also known as bovine pancreatic trypsin inhibitor, BPTI) is a basic protein present in several bovine organs and tissues, such as the lymph nodes, pancreas, lungs, parotid gland, spleen and liver.

Aprotinin is known to inhibit various serine proteases, including trypsin, chymotrypsin, plasmin and kallikrein, and is used therapeutically in the treatment of acute pancreatitis, various stages of shock syndrome, hyperfibrinolytic hemorrhage and myocardial infarction. In addition, administration of aprotinin in high doses significantly reduces blood loss in connection with cardiac surgery, including cardiopulmonary bypass (Bidstrup, *et al.* (1989) *Cardiovasc Surg.* 44:640-645)

As demonstrated in more detail in the examples that follow, plastid expression constructs are employed to direct the expression of aprotinin and a human growth hormone from the plant plastid.

Other sequences which may find use in the production of human biologics include sequences encoding for insulin or insulin precursors may find use in the expression constructs of the present invention. The skilled artisan will recognize that many nucleotide sequences encoding for human biologics may be employed in the constructs of the present invention to direct their expression from a plant plastid such as those described in Goodman and Gelman (1990) *Pharmacological Basis of Therapeutics*, Pergaman Press, 8<sup>th</sup> Edition, Sections 14 and 15.

In developing the constructs the various fragments comprising the regulatory regions and open reading frame may be subjected to different processing conditions, such as ligation, restriction enzyme digestion, PCR, *in vitro* mutagenesis, linkers and adapters addition, and the like. Thus, nucleotide transitions, transversions, insertions, deletions, or the like, may be performed on the DNA which is employed in the regulatory regions or the

DNA sequences of interest for expression in the plastids. Methods for restriction digests, Klenow blunt end treatments, ligations, and the like are well known to those in the art and are described, for example, by Maniatis *et al.* (in *Molecular cloning: a laboratory manual* (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

5           During the preparation of the constructs, the various fragments of DNA will often be cloned in an appropriate cloning vector, which allows for amplification of the DNA, modification of the DNA or manipulation of the DNA by joining or removing sequences, linkers, or the like. Preferably, the vectors will be capable of replication to at least a relatively high copy number in *E. coli*. A number of vectors are readily available for  
10 cloning, including such vectors as pBR322, vectors of the pUC series, the M13 series vectors, and pBluescript vectors (Stratagene; La Jolla, CA).

          In order to provide a means of selecting the desired plant cells, vectors for plastid transformation typically contain a construct which provides for expression of a selectable marker gene. Marker genes are plant-expressible DNA sequences which express a  
15 polypeptide which resists a natural inhibition by, attenuates, or inactivates a selective substance, *i.e.*, antibiotic, herbicide *etc.*.

          Alternatively, a marker gene may provide some other visibly reactive response, *i.e.*, may cause a distinctive appearance or growth pattern relative to plants or plant cells not expressing the selectable marker gene in the presence of some substance, either as applied  
20 directly to the plant or plant cells or as present in the plant or plant cell growth media.

          In either case, the plants or plant cells containing such selectable marker genes will have a distinctive phenotype for purposes of identification, *i.e.*, they will be distinguishable from non-transformed cells. The characteristic phenotype allows the identification of cells, cell groups, tissues, organs, plant parts or whole plants containing  
25 the construct.

          Detection of the marker phenotype makes possible the selection of cells having a second gene to which the marker gene has been linked. This second gene typically comprises a desirable phenotype which is not readily identifiable in transformed cells, but

which is present when the plant cell or derivative thereof is grown to maturity, even under conditions wherein the selectable marker phenotype itself is not apparent.

The use of such a marker for identification of plant cells containing a plastid construct has been described by Svab *et al.* (1993, *supra*). In the examples provided  
5 below, a bacterial *aadA* gene is expressed as the marker under the regulatory control of chloroplast 5' promoter and 3' transcription termination regions, specifically the regulatory regions of the *psbA* gene (described in Staub *et al.*, *EMBO J.*(1993) 12(2):601-606). Numerous additional promoter regions may also be used to drive expression of the selectable marker gene, including various plastid promoters and bacterial promoters which  
10 have been shown to function in plant plastids.

Expression of the *aadA* gene confers resistance to spectinomycin and streptomycin, and thus allows for the identification of plant cells expressing this marker. The *aadA* gene product allows for continued growth and greening of cells whose chloroplasts comprise the selectable marker gene product. Cells which do not contain the selectable marker gene  
15 product are bleached. Selection for the *aadA* gene marker is thus based on identification of plant cells which are not bleached by the presence of streptomycin, or more preferably spectinomycin, in the plant growth medium.

A number of markers have been developed for use with plant cells, such as resistance to chloramphenicol, the aminoglycoside G418, hygromycin, or the like. Other  
20 genes which encode a product involved in chloroplast metabolism may also be used as selectable markers. For example, genes which provide resistance to plant herbicides such as glyphosate, bromoxynil or imidazolinone may find particular use. Such genes have been reported (Stalker *et al.*, *J. Biol. Chem.* (1985) 260:4724-4728 (glyphosate resistant EPSP); Stalker *et al.*, *J. Biol. Chem.* (1985) 263:6310-6314 (bromoxynil resistant nitrilase  
25 gene); and Sathasivan *et al.*, *Nucl. Acids Res.* (1990) 18:2188 (AHAS imidazolinone resistance gene)).

Stable transformation of tobacco plastid genomes by particle bombardment is reported (Svab *et.al.* (1990), *supra*) and Svab *et al.* (1993), *supra*). The methods described therein may be employed to obtain plants homoplasmic for plastid expression constructs.



Generally, bombarded tissue is cultured for approximately 2 days on a cell division-promoting media, after which the plant tissue is transferred to a selective media containing an inhibitory amount of the particular selective agent, as well as the particular hormones and other substances necessary to obtain regeneration for that particular plant species. Shoots are then subcultured on the same selective media to ensure production and selection of homoplasmic shoots.

Transplastomic tobacco plants are analyzed for a pure population of transformed plastid genomes (homoplasmic lines). Homoplasmy is verified using Southern analysis employing nucleic acid probes spanning a region of the transgene and chloroplast genome (i.e. the insertion region). Transplastomic plants which are heteroplasmic (i.e. contain a mixture of plastid genomes containing and lacking the transgene) are characterized by a hybridization pattern of wild type and transgenic bands. Homoplasmic plants show a hybridization pattern lacking the wild type band.

Alternatively, homoplasmy may be verified using the polymerase chain reaction (PCR). PCR primers are utilized which are targeted to amplify from sequences from the insertion region. For example, a pair of primers may be utilized in a PCR reaction. One primer amplifies from a region in the transgene, while the second primer amplifies from a region proximal to the insertion region towards the insertion region. A second PCR reaction is performed using primers designed to amplify the region of insertion. Transplastomic lines identified as homoplasmic produce the expected size fragment in the first reaction, while they do not produce the predicted size fragment in the second reaction.

Where transformation and regeneration methods have been adapted for a given plant species, either by *Agrobacterium*-mediated transformation, bombardment or some other method, the established techniques may be modified for use in selection and regeneration methods to produce plastid-transformed plants. For example, the methods described herein for tobacco are readily adaptable to other solanaceous species, such as tomato, petunia and potato.

For transformation of soybean, particle bombardment as well as *Agrobacterium*-mediated nuclear transformation and regeneration protocols have been described (Hinchey

*et al.* USPN 5,416,011, and Christou *et al.* USPN 5,015,580). The skilled artisan will recognize that protocols described for soybean transformation may be used

In *Brassica*, *Agrobacterium*-mediated transformation and regeneration protocols generally involve the use of hypocotyl tissue, a non-green tissue which might contain a low  
5 plastid content. Thus, for *Brassica*, preferred target tissues would include microspore-derived hypocotyl or cotyledonary tissues (which are green and thus contain numerous plastids) or leaf tissue explants. While the regeneration rates from such tissues may be low, positional effects, such as seen with *Agrobacterium*-mediated transformation, are not expected, thus it would not be necessary to screen numerous successfully transformed  
10 plants in order to obtain a desired phenotype.

For cotton, transformation of *Gossypium hirsutum* L. cotyledons by co-cultivation with *Agrobacterium tumefaciens* has been described by Firoozabady *et al.*, *Plant Mol. Bio.* (1987) 10:105-116 and Umbeck *et al.*, *Bio/Technology* (1987) 5:263-266. Again, as for *Brassica*, this tissue may contain insufficient plastid content for chloroplast  
15 transformation. Thus, as for *Brassica*, an alternative method for transformation and regeneration of alternative target tissue containing chloroplasts may be desirable, for instance targeting green embryogenic tissue.

Other plant species may be similarly transformed using related techniques. Alternatively, microprojectile bombardment methods, such as described by Klein *et al.*  
20 (*Bio/Technology* 10:286-291) may also be used to obtain nuclear transformed plants comprising the viral single subunit RNA polymerase expression constructs described herein. Cotton transformation by particle bombardment is reported in WO 92/15675, published September 17, 1992. Suitable plants for the practice of the present invention include, but are not limited to, soybean, cotton, alfalfa, oil seed rape, flax, tomato, sugar  
25 beet, sunflower, potato, tobacco, *maize*, wheat, rice and lettuce.

The vectors for use in plastid transformation preferably include means for providing a stable transfer of the plastid expression construct and selectable marker construct into the plastid genome. This is most conveniently provided by regions of homology to the target plastid genome. The regions of homology flank the construct to be

transferred and provide for transfer to the plastid genome by homologous recombination, via a double crossover into the genome. The complete DNA sequence of the plastid genome of tobacco has been reported (Shinozaki *et al.*, *EMBO J.* (1986) 5:2043-2049). Complete DNA sequences of the plastid genomes from liverwort (Ohyama *et al.*, *Nature* 5 (1986) 322:572-574) and rice (Hiratsuka *et al.*, *Mol. Gen. Genet.* (1989) 217:185-194), have also been reported.

Where the regions of homology are present in the inverted repeat regions of the plastid genome (known as IRA and IRB), two copies of the transgene are expected per transformed plastid. Where the regions of homology are present outside the inverted repeat regions of the plastid genome, one copy of the transgene is expected per transformed plastid. The regions of homology within the plastid genome are approximately 1kb in size. Smaller regions of homology may also be used, and as little as 100 bp can provide for homologous recombination into the plastid genome. However, the frequency of recombination and thus the frequency of obtaining plants having transformed plastids decreases with decreasing size of the homology regions.

Examples of constructs having regions of homology the plastid genome are described in Svab *et.al.* (1990 *supra*), Svab *et al.* (1993 *supra*) and Zoubenko *et al.* (*Nuc Acid Res* (1994) 22(19):3819-3824).

As described in more detail in the examples below, constructs are described which provide for enhanced expression of DNA sequences in plant plastids. Various promoter/ribosome binding site sequences are employed to direct expression in plant plastids. Promoter sequences of the 16S ribosomal RNA operon (*Prrn*) are linked to a ribosome binding site (RBS) derived from the T7 bacteriophage gene 10 leader sequence (G10L). DNA sequences expressed under the regulatory control of the *Prrn*/G10L sequence show a significantly higher level of protein expression than those levels obtained under the control of other promoter/RBS combinations, while expression of mRNA may or may not be higher in these plants.

In the examples below, nucleic acid sequences encoding CP4 EPSP synthase (USPN 5,633,435) are placed into expression constructs for expression of EPSP synthase

enzyme from the plant plastid. Furthermore, a DNA sequence encoding for hGH (USPN 5,424,199) is also placed into expression construct for the expression of human growth hormone from the plant plastid. The constructs prepared utilize a ribosome binding site designed after the T7 bacteriophage gene 10 leader (G10L) to increase the expression of  
5 the nucleic acid sequences in the plant plastid.

Plastid expression constructs encoding for the expression of EPSPS and hGH are introduced via a chloroplast transformation vector.

Tobacco lines containing the native encoding sequence to the EPSPS enzyme expressed in plastids under the control of the *Prrn*/G10L promoter/ribosome binding site  
10 sequence demonstrate a significantly higher level of protein expression than those levels obtained from EPSPS expressed under the control of the *Prrn*/rbcL RBS sequence. However, EPSPS mRNA is expressed at a higher level in plants expressing CP4 EPSPS from the plastid under the control of the *Prrn*/rbcL(RBS). These results indicate that translation from transcripts containing the T7 bacteriophage gene 10 ribosome binding site  
15 is more efficient. In addition, protein expression levels of EPSPS obtained from transplastomic tobacco lines expressing EPSPS under the control of the *Prrn*/G10L RBS provide for a high level of glyphosate tolerance.

Furthermore, transplastomic tobacco lines transformed to express hGH under the control of the *Prrn*/G10L promoter/ribosome binding site sequence demonstrate a  
20 significantly higher level of protein expression than those levels obtained from hGH expressed under the control of the *PpsbA* promoter/RBS sequence.

Increases in protein expression levels of at least approximately 200 fold may be obtained from constructs utilizing *Prrn*/G10L ribosome binding site for expression of EPSPS and hGH over the expression levels obtained from other promoter/RBS  
25 combinations for plastid expression. In addition, protein levels obtained from plastid expression constructs utilizing the *Prrn*/G10L promoter/RBS sequence may accumulate 50 to 3500 fold higher levels than from nuclear expression constructs. Thus, inclusion of the G10L ribosome binding site in plastid expression constructs may find use for increasing the levels of protein expression from plant plastids.

Thus, the constructs and methods of the present invention provide a means for obtaining transplastomic plants with high level tolerance of herbicides. High levels of tolerance include tolerance of vegetative tissue when amounts of greater than about 16 oz/acre glyphosate are applied, preferably greater than about 32 oz/acre, more preferably greater than about 64 oz/acre, most preferably greater than about 128 oz/acre. Furthermore, high levels of tolerance can also include tolerance of reproductive tissues when amounts of greater than about 16 oz/acre glyphosate are applied, preferably greater than about 32 oz/acre, most preferably greater than about 64 oz/acre.

The invention now being generally described, it will be more readily understood by reference to the following examples which are included for purposes of illustration only and are not intended to limit the present invention.

## EXAMPLES

### Example 1 Expression Constructs

Constructs and methods for use in transforming the plastids of higher plants are described in Zoubenko *et al.* (*Nuc Acid Res* (1994) 22(19):3819-3824), Svab *et al.* (*Proc. Natl. Acad. Sci.*(1990) 87:8526-8530 and *Proc. Natl. Acad. Sci.*(1993) 90:913-917) and Staub *et al.* (*EMBO J.* (1993) 12:601-606). Constructs and methods for use in transforming plastids of higher plants to express DNA sequences under the control of a nuclearly encoded, plastid targeted T7 polymerase are described in U.S. Patent Number 5,576,198. The complete DNA sequences of the plastid genome of tobacco are reported

by Shinozaki *et al.* (*EMBO J.* (1986) 5:2043-2049). All plastid DNA references in the following description are to the nucleotide number from tobacco.

The complete nucleotide sequence encoding the tobacco cytochrome *f* (*petA*) is described in Bassham *et al.*, (1991) *J Biol Chem* 266:23606-23610 and Konishi *et al.*

5 (1993) *Plant Cell Physiol* 34:1081-1087.

#### 1A. Promoter/Ribosome Binding Site Sequences

The promoter region of the plastid 16S ribosomal RNA operon (*Prrn*) is linked to a synthetic ribosome binding site (RBS) patterned on the plastid *rbcL* gene leader to create the *Prrn/rbcLRBS* fragment. The *Prrn/rbcLRBS* sequence is as described in Svab *et al.*

10 (1993, *supra*) for the *Prrn/rbcL*(S) fragment.

The promoter region of the plastid *psbA* promoter (*PpsbA*) and terminator sequences (*TpsbA*) are described in Staub *et al.* (1993, *EMBO J.*, 12, 601-606).

The *Prrn/G10L* sequence was constructed by annealing two oligonucleotide sequences, T7lead1 and T7lead2 (Table 1), to create the G10L plastid ribosome binding site (Figure 1). The G10L sequence was ligated to the 3' terminus of the *Prrn* promoter sequence as an *EcoRI/NcoI* fragment to create the *Prrn/G10L* sequence.

Table 1

---

<b>T7lead1</b>	5'-AAT TGT AGA AAT AAT TTT GTT TAA CTT TAA GAA GGA GAT ATA CC-3'
----------------	---

20

<b>T7lead2</b>	5'-CAT GGG TAT ATC TCC TTC TTA AAG TTA AAC AAA ATT ATT TCT AC-3'
----------------	---

---

Chimeric genes are preferably inserted into the expression vector to direct their transcription from the *Prrn* promoter. Thus, in the plastid genome, chimeric genes are transcribed from the *Prrn*/RBS promoter, or the *Prrn/G10L* promoter in the plant plastid. The nucleic acid sequence of the *Prrn/G10L* fusion is provided in figure 6.

#### 1B. CP4 EPSPS Plastid Expression Constructs

A plastid expression vector pMON30117 is constructed from a precursor vector pPRV111B (Zoubenko, *et al.* 1994, *supra*, GenBank accession U12813). The vector pMON30117 carries a multiple cloning site for insertion of a passenger gene in a *Prm/rbcLRBS/Trps16* expression cassette. The *Prm/rbcLRBS* sequence is cloned into  
5 pPRV111B vector as an *EcoRI/NcoI* fragment, and the terminator region from the plastid *rps16* gene (*Trps16*) is cloned 3' of the *Prm* promoter as a *HindIII/NcoI* fragment. The *Trps16* fragment comprises the *rps16* gene 3'-regulatory region from nucleotides 5,087 to 4,939 in the tobacco plasmid DNA.

The pPRV111B backbone of the vector pMON30117 contains a marker gene,  
10 *aadA*, for selection on spectinomycin and streptomycin, and *rps 7/12* for the integration, by homologous recombination, of the passenger DNA into *tmV-rps7/12* intergenic region.

A nuclear expression construct, pMON10154, was prepared as a control for integration into plants by *Agrobacterium*-mediated transformation. In this construct, the CP4 native gene is expressed from the constitutive Figwort Mosaic Virus promoter and the  
15 *Petunia* HSP70 leader, and has the E9 terminator. Targeting to plastids is by the chloroplast transit peptide of the *Petunia* EPSPS translationally fused to the N-terminus of the CP4 gene.

The plastid expression construct pMON30118 was prepared by cloning the native CP4 EPSPS gene fused with the N-terminal five (5) amino acids from the plastid *rbcL*  
20 (described in Svab *et al.*, 1993 *supra*) gene as an *NcoI/SmaI* fragment into the multiple cloning site of the vector pMON30117.

The plastid expression construct pMON30123 is essentially the same as pMON30118 with the exception of the deletion of the N-terminal five (5) amino acids from the plastid *rbcL*.

25 The plastid expression construct pMON30130 was created by replacing the native CP4 EPSPS of pMON30123, with a synthetic CP4 gene. This construct also lacks the N-terminal 5 amino acid fusion from the plastid *rbcL* gene.

The plastid expression construct pMON38773 was constructed by replacing the *Prm*/RBS sequence of pMON30123 with the *Prm*/G10L promoter sequence described

above. The EPSPS DNA sequence of pMON38773 also lacks the N-terminal 5 amino acid fusion from the plastid *rbcL* gene.

A plastid expression construct, pMON38766 was constructed using the promoter from T7 phage gene 10 (P-T7), including G10L, CP4 (native) gene coding region, and the  
5 terminator sequence from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON38797 was constructed using the promoter from T7 phage gene 10 (P-T7), including G10L, CP4 (synthetic) gene coding region, terminator from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON38798 was constructed using the promoter  
10 of the 16SrDNA operon (Prn), G10L, CP4 (synthetic) gene coding region, terminator from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON38793 was constructed using the promoter of the 16SrDNA operon (Prn), a synthetic ribosome binding site (RBS) patterned from the plastid *rbcL* gene, the glyphosate tolerant *Petunia* EPSP synthase gene (P-EPSPS,  
15 Padgett, *et al.* (1987) *Arch. Biochem. Biophys.* 258:564-573) carrying the mutation Glycine to Alanine at amino acid position 101, terminator from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON38796 was constructed using the promoter of the 16SrDNA operon (Prn), synthetic ribosome binding site (RBS) patterned from the  
20 plastid *rbcL* gene, the glyphosate tolerant *Achromobacter* (strain LBAA) EPSP synthase gene (U.S. Patent Number 5,627,061, the entirety of which is incorporated herein by reference) carrying the mutation Glycine to Alanine at amino acid position 100 (G100A), terminator from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON45204, was constructed using the promoter  
25 of the 16SrDNA operon (Prn) with the G10L, the glyphosate tolerant *Pseudomonas* (strain LBAA) EPSP synthase gene carrying the mutation Glycine to Alanine at amino acid position 100 (G100A), terminator from plastid *rps16* gene (Trps16).

A plastid expression construct, pMON45201, was constructed using the promoter of the 16SrDNA operon (Prn), synthetic ribosome binding site (RBS) patterned from the



plastid *rbcl* gene, wild-type glyphosate tolerant *Bacillus subtilis* *aroE* (EPSPS)(U.S. Patent Number 5,627,061) gene, terminator from plastid *rps16* gene (*Trps16*).

A plastid expression construct, pMON45259, was constructed using the promoter of the 16SrDNA operon (*Prrn*) with the G10L sequence functionally associated with the  
5 nucleic acid sequence encoding the synthetic CP4 protein having an additional sequence at the N-terminus encoding the first 14 amino acids of the green fluorescent protein (GFP) (GKGEELFTGVVPSM). The sequence encoding the 14 amino acid GFP fusion begins at the glycine in the second position of the protein. The construct also contains the *rps16* terminator.

10 Another plastid expression construct, pMON49218, was constructed to express the synthetic CP4 sequence with the 14 amino acid GFP fusion from the promoter region of the 16SrDNA operon having the nuclear-encoded RNA polymerase region (*Prrn*PEP+NEP), and the terminator region from the plastid *rps16* gene. The DNA  
15 sequence of the *Prrn*/NEP/G10L::14aaGFP fusion is provided in figure 7.

#### 1C. Bucril (*bxn*) Plastid Expression Constructs

The *bxn* herbicide resistance gene (U.S. Patent Number 4,810,648, the entirety of which is incorporated herein by reference) was removed from the plasmid pBrx47 as an *Nco* I to *Asp*718 restriction fragment and cloned into *Nco*/*Asp*718 cut pUC120 resulting in  
20 plasmid pBrx87. Plasmid pBrx87 was then digested with *Nco*/*Xba* and cloned into the *Nco*/*Xba* sites of the plasmid pLAA21 which contains the *Prrn* plastid promoter and the *rpsL* 3' region for plastid expression. The resulting plasmid was designated pBrx89. Plasmid pBrx89 was digested with *Sac* I and *Hind* III and the 1.5 kb chimeric *bxn* gene with plastid expression signals was inserted into the *Sac*/*Hind* III sites of the tobacco  
25 plastid homology vector pOVZ44B (Zoubenko et al, Nuc Acids Res 22: 3819-3824 (1994)) to create plasmid pCGN5175.

To construct plasmid pCGN6114, plasmid pBrx90 (a Bluescript plasmid containing the *bxn* gene encoding the bromoxynil specific nitrilase) was digested with *Nco* I/*Asc* I and the *bxn* structural gene was substituted for the GUS gene in the *Nco*/*Asc*

digested plasmid pCGN5063 resulting in plasmid pCGN6107. This plasmid contains the the *bxn* gene under the control of the T7 promoter/gene10 leader at the 5' end and the *psbA*/T7 hybrid transcriptional terminator at the 3' end of the chimeric gene. This T7 promoter/*bxn* chimeric gene was excised from pCGN6107 as a *Hind* III/*Not* I DNA  
5 segment and moved into the chloromphenical plasmid BCSK+ (Stratagene) at the *Hind* III/*Not* sites to create plasmid pCGN6109. The chimeric gene was then moved as a *Hind* III/*Not* fragment from pCGN6109 into the chloroplast homology vector pOVZ44B described above to create plasmid pCGN6114. Tobacco plants transformed with pCGN6114 require the T7 RNA polymerase be provided in the plant plastid background to  
10 activate transcription of the chimeric *bxn* gene via the T7 promoter. This system has previously been detailed in McBride et al PNAS 91:7301-7305 (1994) and McBride et al US patent number 5,576,198.

#### 1D. BXN/AHAS Plastid Expression Constructs

15 A plastid expression construct, pCGN5026, is prepared to direct the expression of BXN and AHAS from the plant plastid. The AHAS nucleotide sequence (described in EP Publication Number 0 525 384 A2, the entirety of which is incorporated herein by reference) is translationally linked to the BXN nucleotide sequence (U.S. Patent Number 4,810,648, the entirety of which is incorporated herein by reference). The AHAS structural  
20 gene encoding acetohydroxyacid synthase was cloned from the plasmid pCGN4277 as an *Nco* I to *Age* DNA fragment into the *Nco*/*Xma* sites of plasmid pUC120 to create plasmid pCGN5022. This plasmid was then digested with the enzymes *Bam*H I and *Pst* and a 1.3 kb *Bam*/*Pst* DNA segment containing the *bxn* gene encoding the bromoxynil-specific nitrilase was excised from the plasmid pBrx26 and cloned into the *Bam*/*Pst* sites of  
25 pCGN5022 to create plasmid pCGN5023. Plasmid pCGN5023 contained a 3.3 kb DNA segment containing the AHAS/*bxn* operon segment and this fragment. This plasmid was cut at the unique *Pst* site and this *Pst* site was removed and replaced with a synthetic linker containing a unique *Xba* I restriction site generating plasmid pCGN5024. Plasmid pCGN5024 was digested with *Nco*/*Xba* and the 3.3 kb *Nco*/*Xba* DNA fragment was

cloned into the plastid promoter cassette vector pLAA21(Pst) that had been digested with *Nco* and *Xba* to remove the GUS gene. The plasmid resulting from this cloning was designated plasmid pCGN5025 and contained the herbicide operon under the control of the plastid promoter *Prrn* and the *rpsL* 3' DNA segment. The entire chimeric herbicide operon under the control of the plastid expression elements was excised from pCGN5025 as a *Sac* I/*Pst* DNA fragment and cloned into the *Sac*/*Pst* sites of the plastid homology cassette vector pOVZ44B (Zoubenko *et al*, *Nuc Acids Res* 22:3819-3824 (1994)) to facilitate transfer into the tobacco chloroplast genome.

#### 10 1E. Bt *cryI*Ac and *bxn* Plastid Expression Construct

Plasmid pBrx9 (Stalker and McBride, (1987) *J Bacteriol* 169:955-960), an original clone from *Klebsiella* containing a *bxn* gene DNA segment, was used as a template to generate an ~450 bp *Bam*H I/*Cla* I PCR DNA fragment that encompasses the N-terminal end of the *bxn* gene and includes 44 bp of the 5' untranslated portion of the native gene. This fragment was exchanged with the ~400 bp *Bam*/C*la* fragment in the plasmid pBrx90 resulting in plasmid pBrx90.1. This plasmid contains the entire *bxn* gene and the 44 bp untranslated 5' DNA segment. The *bxn* gene was excised from plasmid pBrx90.1 as a *Bam*/A*sc* I DNA segment and inserted into plasmid pCGN5146 at the *Bgl* II/A*sc* I sites to generate plasmid pCGN5191. Plasmid pCGN5146 is a pKK233-2 (Pharmacia) derivative containing the full-length *cryI*Ac gene encoding the HD-73 Bt protoxin. Plasmid pCGN5191 then contains the *cryI*Ac and *bxn* genes in an operon configuration with the *bxn* gene being the distal gene in the operon. Both genes are under the control of the *Ptac* promoter for *E coli* expression in 5191. Plasmid pCGN5191 was digested with *Nco*/A*sc* and the *Nco*/A*sc* DNA fragment containing the Bt/*bxn* operon was cloned into the *Nco*/A*sc* sites of the chloroplast homology vector pCGN5155, a derivative of pOVZ44B. The resulting plasmid, pCGN5197 contains the Bt/*bxn* operon under the control of the *Prrn* plastid promoter and *rpsL* transcription terminator regions. This plasmid facilitated transfer of the Bt/*bxn* chimeric operon into the tobacco plastid genome.

#### 1F. Phytoene desaturase Plastid Expression Constructs

The *crtI* gene was obtained as a *Hind* III/*Sal* I PCR fragment from the original plasmid containing the *Erwinia carotova crt* operon (Misawa et al, (1994) *Plant Jour* 6:481-489)) and cloned as a *Hind* III/*Sal* I DNA segment into BCSK+ (Stratagene) at the

5 *Hind* III/*Sal* I sites to generate plasmid pCGN5172. The *crtI* fragment was cloned from pCGN5172 as an *Nco* I/*Sal* I fragment into pCGN5038 (a derivative of pOVZ44B) to create the plastid expression construct pCGN5177. This construct directs the expression of the *crtI* sequence from the *Prm* promoter and the *rps16* terminator sequence. This plasmid facilitated the transfer of the chimeric *crtI* gene into the tobacco plastid genome.

10

#### 1G. hGH Expression Constructs for Plant Transformation

##### Nuclear Expression Constructs

The construct pWRG4747 was constructed to direct the expression of hGH in the plant nuclear genome. This vector contains the hGH operably linked to the Figwort

15 Mosaic Virus promoter (USPN 5,378,619, the entirety is incorporated herein by reference) and the CTP2 leader for directing the hGH protein into the plastid. The FMV/CTP2L::hGH::NpA fragment is cloned along with the DNA sequence conferring resistance to Kanamycin between the right and left borders (RB and LB) of the transfer DNA (tDNA) of *Agrobacterium tumefaciens* to direct the integration into the nuclear

20 genome.

The nuclear transformation vector pWRG4744 contains essentially the same elements as pWRG4747 except the construct lacks the CTP2 leader and the hGH protein is directed to the plant cell cytoplasm.

##### Plastid Expression Constructs

25 The plastid expression vector pWRG4838 was constructed using the full length hGH gene expressed from the promoter region from the *psbA* gene and the *psbA* gene terminator, *PpsbA* and *TpsbA* respectively (described in Staub *et al.* (1993), *supra*). This chimeric promoter-gene-terminator fusion (*PpsbA*::hGH:: *TpsbA*) is cloned adjacent to the selectable marker gene *aadA* also driven by the plastid expression elements of the *psbA*

gene. The two chimeric gene sequences are cloned into a vector between two sequences which direct the integration of the chimeric gene sequences into the tobacco plastid genome upstream of the plastid 16SrDNA. This is joined to a 1 kb Ampicillin resistance gene which provides for selection of *E. coli* containing the construct and the pUC origin of replication for plasmid maintenance in *E. coli*.

The plastid expression construct pMON38755 was prepared using the hGH DNA sequence translationally fused at the N-terminus with the yeast ubiquitin gene, creating the Ubi-hGH fusion gene. The Ubi-hGH fusion gene is cloned next to the *aadA* gene for selection of transplastomic tobacco on media containing spectinomycin or streptomycin (from pPRV112B described in Zoubenko *et al.* (1994) *supra*). Sequences are included for the homologous recombination of sequences encoding for hGH and *aadA* expression. These sequences are obtained from the vector pPRV112B described in Zoubenko *et al.* (1994, *supra*). This is joined to a 1 kb Ampicillin resistance gene which provides for selection of *E. coli* containing the construct and the pUC origin of replication for plasmid maintenance in *E. coli*.

The plastid expression construct pMON38794 contains essentially the same elements as pMON38755, with the following exception. The 0.15 kb *psbA* promoter sequence is replaced with the *Prn/G10L* promoter sequence described above.

## 1H. Constructs for the Expression of Aprotinin in Plastids

A series of constructs were prepared to direct the expression of the pharmaceutical protein aprotinin from the plastid. The nucleic acid sequence encoding for aprotinin (Figure 2) was cloned into a plastid expression construct to control the expression of aprotinin from the T7 gene 10 leader promoter which is induced from a nuclearly expressed, plastid targeted T7 Polymerase. The constructs used in which the aprotinin sequence was cloned are as described in U.S. Patent Number 5,576,198, the entirety of which is incorporated herein by reference. The plastid transformation vector pCGN6146 is designed by replacing the DNA sequence encoding for GUS from pCGN4276 (described in USPN 5,576,198) with the coding sequence of aprotinin. The tobacco plastid

transformation construct pCGN6147 contains the same elements as pCGN6146 except pCGN6147 contains the six 5' amino acids of the GUS encoding sequence ligated to the 5' terminus of the aprotinin encoding sequence. The six amino acids of the 5' terminus of the GUS nucleotide sequence are included to aid in the translation of the aprotinin protein.

5 The tobacco plastid transformation vector pCGN6156 is essentially the same as pCGN4276 except the coding region of aprotinin is cloned to the 3' end of the GUS coding sequence. Thus, pCGN6156 contains as operably linked the T7 promoter, a DNA sequence encoding for GUS fused with the DNA sequence encoding for aprotinin and the psbA 3' transcription termination sequence.

10 A plastid expression construct, pCGN6154, was constructed from pCGN4276 by replacing the GUS coding sequence with the aprotinin protein operably linked to the 3' terminus of the coding sequence of cytochrome *f* (*petA*) of the tobacco chloroplast. Thus, pCGN6154 contains the T7 promoter sequence operably linked to the nucleotide sequence of *petA* and aprotinin. The *petA* sequence is included to direct the expressed aprotinin  
15 protein to the thylakoid.

## Example 2 Plant Transformation

### 2A. Nuclear Transformation

Tobacco plants transformed to express the constructs pWRG4744 and pWRG4747  
20 in the nucleus of a plant cell may be obtained as described by Horsch *et al.* (*Science* (1985) 227:1229-1232).

### 2B. Plastid Transformation

Tobacco plastids are transformed by particle gun delivery of microprojectiles as described by Svab and Maliga (*Proc. Natl. Acad. Sci.* (1993) 90:913-917), and described  
25 here.

Dark green, round leaves are cut, preferably from the middle of the shoots, from 3-6 week old *Nicotiana tabacum* cv. Havana which have been maintained *in vitro* on hormone free MS medium (Murashige and Skoog, (1962) *Physiol Plant.* 15, 473-497) supplemented with B5 vitamins in Phytatrays or sundae cups with a 16 hour photoperiod at

24°C. Each cut leaf is then placed adaxial side up on sterile filter paper over tobacco shoot regeneration medium (TSO medium: MS salts, 1mg/l *N*<sup>6</sup>-benzyladenine, 0.1mg/l 1-naphthaleneacetic acid, 1 mg/l thiamine, 100mg/l inositol, 7g/l agar pH 5.8 and 30g/l sucrose). Leaves are preferably placed in the center of the plate with as much contact with the medium as possible. The plates are preferably prepared immediately prior to use, but may be prepared up to a day before transformation by particle bombardment by wrapping in plastic bags and storing at 24°C overnight.

Tungsten or gold particles are sterilized for use as microcarriers in bombardment experiments. Particles (50mg) are sterilized with 1 ml of 100% ethanol, and stored at -20°C or -80°C. Immediately prior to use, particles are sedimented by centrifugation, washed with 2 to 3 washes of 1 ml sterile deionised distilled water, vortexed and centrifuged between each wash. Washed particles are resuspended in 500 µl 50% glycerol.

Sterilized particles are coated with DNA for transformation. Twenty-five micoliter aliquots of sterilized particles are added to a 1.5 ml microfuge tube, and 5 µg of DNA of interest is added and mix by tapping. Thirty-five microliters of a freshly prepared solution of 1.8M CaCl<sub>2</sub> and 30 mM spermidine is added to the particle/DNA mixture, mixed gently, and incubated at room temperature for 20 minutes. The coated particles are sedimented by centrifuging briefly. The particles are washed twice by adding 200µl 70% ethanol, mixing gently, and centrifuging briefly. The coated particles are resuspended in 50µl of 100% ethanol and mixed gently. Five to ten microliters of coated particles are used for each bombardment.

Transformation by particle bombardment is carried out using the PDS 1000 Helium gun (Bio Rad, Richmond, CA) using a modified protocol described by the manufacturer.

Plates containing the leaf samples are placed on the second shelf from the bottom of the vacuum chamber and bombarded using the 1100 p.s.i. rupture disk. After bombardment, petriplates containing the leaf samples are wrapped in plastic bags and incubated at 24°C for 48 hours.

After incubation, bombarded leaves are cut into approximately 0.5 cm<sup>2</sup> pieces and placed abaxial side up on TSO medium supplemented with 500 µg/ml spectinomycin.

After 3 to 4 weeks on the selection medium, small, green spectinomycin resistant shoots will appear on the leaf tissue. These shoots will continue to grow on spectinomycin containing medium and are referred to as primary putative transformants.

When the primary putative transformants have developed 2 to 3 leaves, 2 small pieces (approximately 0.5 cm<sup>2</sup>) are cut from each leaf and used for either selection or for a second round of shoot regeneration. One piece is placed abaxial side up on plates containing TSO medium supplemented with 500 µg/ml spectinomycin, and the other piece is placed abaxial side up on TSO medium supplemented with 500 µg/ml each of spectinomycin and streptomycin. Positive transformants are identified as the shoots which form green callus on the TSO medium containing spectinomycin and streptomycin.

After 3 to 4 weeks, the tissue placed on TSO medium containing only spectinomycin, which has been identified as positive on the TSO medium with spectinomycin and streptomycin, will develop green shoots. Two to four shoots of each positive transformant are selected and transferred to TSO medium supplemented with 500 µg/ml spectinomycin for generation of roots. Southern analysis is performed on 2 shoots to confirm homoplasmy as described below. Shoots from homoplasmic events are transferred to the greenhouse for seed production, while transformants which are not homoplasmic are sent through a second round of regeneration on TSO medium with 500 µg/ml spectinomycin to attain homoplasmy.

20

### **Example 3 Analysis of Transplastomic Tobacco Plants Transformed with Herbicide Tolerance Constructs**

#### **3A. Southern Analysis**

Transformed plants selected for marker *aadA* marker gene expression are analyzed to determine whether the entire plastid content of the plant has been transformed (homoplastic transformants). Typically, following two rounds of shoot formation and spectinomycin selection, approximately 50% of the transgenic plantlets which are analyzed are homoplastic, as determined by Southern blot analysis of plastid DNA. Homoplasmic plantlets are selected for further cultivation.



Genomic DNA is isolated from transformed tobacco plants, electrophoresed, and transferred to filters as described in Svab *et al.* ((1993), *Proc Natl Acad Sci*, 90:913-917).

Homoplasmic tobacco plants transformed to express CP4 EPSPS in plastids were identified using a probe prepared from a 2.4 kb *EcoRI/EcoRV* fragment from the vector pOVZ2 (similar to pOVZ15 described in Zoubenko, *et al.* 1994, *supra*). The 2.4 kb probe  
5 fragment encompasses part of the targeting sequence.

Results of the Southern hybridizations identified 3 homoplasmic lines from tobacco transformed with the constructs pMON30123 and pMON30130 and 1 line from tobacco transformed with pMON38773 for further analysis.

10 The complete disappearance of the 3.27 Kb native tobacco *Bam*HI fragment in the lines 30123-19-1A, 30123-23-2A, 30123-18-1B, 30130-51-2A, 30130-51-2P, 30130-57-1P, and 38773-6 with a probe covering the region of integration, and the appearance of expected sized bands for the inserted DNA fragments in those transformants, 5.14 kb and 0.9 kb, establishes that the transformed plants are homoplasmic for the intended  
15 constructs.

Results of the Southern hybridizations identified 3 homoplasmic lines from tobacco transformed with pCGN5177, lines 74-1B-P, 74-2 and 74-7.

Transplastomic 5175 and 6114 tobacco lines were analyzed by Southern hybridization for homoplasmy as described above. Results of the Southern hybridizations  
20 identified 4 homoplasmic lines from tobacco transformed with pCGN6114.

Results from hybridizations of 5175 transplastomic tobacco lines identified one line, 76-4A-F, as homoplasmic, and a second line as 95% homoplasmic.

Homoplasmic tobacco plants transformed to express BXN/AHAS in plastids were identified using Southern hybridizations as described above.

25 Results of the Southern hybridizations identified 14 homoplasmic lines from tobacco transformed with pCGN5026. The filters were reprobbed with a BXN gene fragment, and 21 lines were found to contain BXN, 14 lines of which were homoplasmic.

### 3B. Northern Analysis

In order to determine the level of transcription of the EPSPS, BXN or AHAS mRNA expressed in the transplastomic tobacco plants, Northern blot hybridizations were performed with total RNA isolated from each of the lines identified. Total RNA was isolated using TRIzol reagent (Gibco-BRL Life Technologies, Gaithersburg, MD) according to the manufacturers protocol. Total RNA, 2µg, was separated on a denaturing agarose gel and transferred to nylon membrane (Maniatis *et al.*, 1989, *supra*). Radioactive probes for hybridizations were prepared using random primer labeled (using Random Primer labeling kit from Boehringer Mannheim) CP4 EPSPS, phytoene desaturase, BXN, or AHAS fragments and hybridizations were carried out in 2x SSPE (Maniatis, *et al.*, 1989, *supra*), at 60°C. Filters were stripped and reprobed with a plastid 16S ribosomal RNA gene probe (from pPRV112A, Zoubenko, *et al.*, 1994, *supra*) to confirm homogenous loading of RNA on the filter.

Results of the Northern hybridizations performed with EPSPS probes demonstrate that all seven (7) lines examined express CP4 EPSPS mRNA. Hybridizations performed with the 16S ribosome probe confirm that denaturing gels were loaded with similar amounts of total RNA for each sample. Furthermore, transplastomic tobacco lines expressing EPSPS from the *Prrn/rbcL*(RBS) (pMON30123) regulatory elements express EPSPS mRNA to higher levels than tobacco plants homoplasmic for EPSPS controlled by the *Prrn/G10L* (pMON38773) promoter/RBS sequences.

Results of Northern hybridizations performed with BXN, AHAS and *crtI* probes demonstrates that all homoplasmic 5026, 5175, and 5177 tobacco lines expressed *crtI*, BXN and/or AHAS mRNA.

### 3C. Western Blot Analysis of Tobacco CP4 EPSPS

To determine the expression of the EPSPS Western blot analysis was performed on a single line from each construct, pMON30123, pMON30130, and pMON38773.

Total soluble protein was extracted from frozen leaf tissue by grinding 250 mg tissue in 250µl of PBS buffer (1 mM KH<sub>2</sub>PO<sub>4</sub>, Na<sub>2</sub>HPO<sub>4</sub>, 0.137M NaCl, 2.7 mM KCl pH 7.0) containing protease inhibitors. The homogenate is centrifuged for 5 minutes, and the

supernatant is transferred to a fresh tube. The concentration of the protein in the supernatant is determined using a protein concentration assay (BioRad, Richmond, CA).

Extracted total protein is electrophoresed on a 4-20% SDS-PAGE gel (Sigma, St Louis, MO), and transferred to PVDF membrane in 1x SDS-PAGE buffer (Maniatis *et al.* 1989, Cold Spring Harbor Press). Standards of quantitated purified CP4 EPSPS protein were used to quantify the expression of the CP4 EPSPS as expressed in the plant plastid.

Western hybridizations are performed as described in Staub and Maliga (1993) *EMBO Journal*, 12(2) 601-606, except using antibodies raised to EPSPS. PVDF membranes containing the transferred electrophoresed protein were incubated in a blocking solution of PBS buffer containing 0.05% Tween-20 (PBS-T) and 5% milk overnight at 4°C. The membranes are then incubated in a solution of PBS-T containing 1% milk and a primary antibody raised in goats to the CP4 EPSPS for 2 hours at room temperature. The membranes are washed three times in a solution of PBS-T containing 0.1% milk, each wash for 5 minutes at room temperature. The membranes are then incubated in a solution of PBS-T containing 1% milk and sheep anti-goat antibody for 1 hour at room temperature, and washed again in PBS-T containing 0.1% milk, three times for 10 minutes at room temperature. A final wash using only PBS-T is performed before developing the membranes using a nonradioactive detection kit (ECL, Amersham).

**Table 2**

Construct Number	% Total Soluble Protein
pMON30123	0.001
pMON30130	0.002
pMON38773	0.2
pMON38798	0.2
pMON45259	>12.0
pMON49218	>12.0

The results listed in Table 2 demonstrate that significant increases in the level of EPSPS protein may be obtained from plants transformed to express EPSPS from the

*Prrm*/G10L promoter. These results demonstrate that EPSPS expression driven by the *Prrm*/rbcLRBS regulatory sequences may produce approximately 0.001% of the total soluble protein as EPSPS, while in plants expressing EPSPS from the *Prrm*/G10L regulatory sequences express 0.2% of the total soluble protein as EPSPS. Subsequent lines  
5 have demonstrated total soluble protein of about 1% EPSPS when expressed from the *Prrm*/G10L regulatory sequences. These results, taken together with the results of the Northern hybridizations above, indicate that more efficient translation may be obtained from the G10L ribosome binding site.

Furthermore, plastid expression constructs containing the N-terminal 14 amino  
10 acid from GFP demonstrated high levels of protein expression. Transplastomic lines containing either pMON45259 or pMON49218 demonstrated total soluble protein of greater than 12% CP4 EPSPS.

Western immunoblot hybridization were also performed on 2 homoplasmic 5026 tobacco lines as described above, using antibodies raised against bromoxynil. The results  
15 of Western immunoblot analysis of total soluble protein extracted from tobacco lines transformed with pCGN5026 demonstrated that both homoplasmic lines produced nitrilase protein.

Western immunoblot analysis was performed as described above from total protein extracted from tobacco lines transformed with pCGN6114 and pCGN5197.

20 The results of the analysis demonstrated that bromoxynil was produced in 6114 tobacco lines ranging from 1% to 2% of the total soluble leaf protein.

The results of the Western analysis of the 20 5197 tobacco lines demonstrated that bromoxynil and Bt were both produced as 1% of the total soluble leaf protein.

### 25 3D. Analysis of EPSPS Enzyme Activity

The EPSPS enzyme activity in transplastomic tobacco plants containing the plastid expression vector pMON38773 was determined using a high pressure liquid chromatography (HPLC) assay.

Methods for the analysis of EPSPS enzyme activity are described in Padgett *et al.* (*J. Biol. Chem.* (1988)263:1798-1802 and *Arch. Biochem. Biophys.* (1987)258:564-573) and Wibbenmeyer *et al.* (*Biochem. Biophys. Res. Commun.* (1988)153:760-766).

The results are summarized in Table 3 below.

5

Table 3

<u>Nuclear</u> Enzymatic Activity Range	<u>Nuclear</u> % Total Plants In Range	<u>Chloroplast</u> 38773-6
1-3.7 $\mu\text{mol/mg}$	1%	
>0.1 $\mu\text{mol/mg}$	16%	
>10 nmol/mg	55%	16.39 nmol/mg
>1 nmol/mg	32%	
0 nmol/mg	3%	

These results demonstrate that EPSPS expression in plastids produces active EPSPS enzyme.

5

### 3E. Analysis for Glyphosate Tolerance

A transplastomic tobacco line homoplasmic for the construct pMON38773 was tested *in vitro* to determine the highest level of glyphosate tolerance. Explant tissue was prepared from leaf pieces of nontransgenic wild type tobacco control, *Havanna*, plants and the homoplasmic tobacco line 38773-6 and cultured for regeneration of shoots on TSO medium (described above) supplemented with glyphosate levels of 50 $\mu\text{M}$ , 75 $\mu\text{M}$ , 100 $\mu\text{M}$ , 150 $\mu\text{M}$  and 200 $\mu\text{M}$ . The results are summarized in Table 4 below. The number of explants producing shoots was determined at 3 weeks and 6 weeks after explant preparation and culturing on glyphosate containing medium.

15

Table 4

Glyphosate Level ( $\mu$ M)	Total Number Explants	Number Regenerating 3 Weeks	Number Regenerating 6 Weeks	% Explant Regeneration
<b>Wild Type</b>				
50	10	0	0	0
75	10	0	0	0
100	10	0	0	0
150	10	0	0	0
200	10	0	0	0
<b>38773-6</b>				
50	8	5	8	100
75	18	14	18	100
100	17	12	15	88
150	18	10	16	89
200	16	8	15	86

The above results demonstrate that at all levels of glyphosate examined, shoots  
 5 regenerated from explants prepared from a tobacco line homoplasmic for pMON38773,  
 while no shoots regenerated from explants prepared from nontransformed control plants.  
 These results suggest that tobacco plants expressing EPSPS in plastids demonstrate  
 tolerance to glyphosate levels of at least 200 $\mu$ M.

Additional transplastomic lines were tested *in vitro* for glyphosate tolerance as  
 10 described above. The results are shown in Table 5

TABLE 5

Summary of tobacco plastid transformation experiments with various constructs containing EPSPS genes.

Construct	Spec/strep (+)	No. of shoots Gly 50 uM(+)
pMON38766 (Wild)	1	0
pMON38766 (T7)	6	0
pMON38773 (Wild)	9	5 (1)
pMON38797 (Wild)	2	0
pMON38798	6	6
pMON38793	8	0
pMON38796	4	0
pMON45201	9	3
pMON45204	12	*

(No. of shoots positive at 1 mM glyphosate)

These results demonstrate that these transplastomic lines show tolerance to glyphosate. The numbers in parentheses are the number of shoots resistant to selection at 1 mM glyphosate. Thus, as can be seen in table 5, tobacco lines are generated that are tolerant of selection at 1 mM glyphosate.

Homoplasmic tobacco plants of the line 38773-6 are sprayed with glyphosate using a track sprayer at concentrations corresponding to 0oz/acre, 16oz/acre, 32oz/acre and 64oz/acre to test for whole plant tolerance. Plant height was measured before and after spraying with glyphosate. The vegetative injury data was collected two weeks after spraying, while the reproductive injury data was collected at plant maturity.

Initial results indicate that homoplasmic tobacco lines sprayed are tolerant of glyphosate at the concentration of 16oz/acre as demonstrated in the vegetative tissue injury (Table 6). As can be seen in Table 5 transplastomic lines were generated which



demonstrated a good level of glyphosate tolerance at 32oz/Acre. In subsequent experiments with additional transformed lines, transplastomic lines have shown tolerance to glyphosate at a level of 64oz/Acre.

5 Tolerance is characterized by the continued growth and greening of tissues sprayed with glyphosate. However, as the concentration of glyphosate applied increased, there was a corresponding increase in the level of vegetative injury. In contrast, nontransformed control plants which were highly susceptible to glyphosate concentrations as low as 16 oz/Acre.

10

Table 6

Plant No.	Construct	Roundup rate (oz/A)	Plant height (cm) before spray	Plant height (cm) after spray	Vegetative injury	Fertility rating
1	38773	0	12.2	30.5	0	0
2	38773	0	13.6	34.0	0	0
3	38773	0	8.6	23.8	0	0
4	38773	0	8.6	26.2	0	0
5	38773	0	7.8	28.8	0	0
6	38773	0	12.8	31.5	0	0
7	38773	0	12.2	31.6	0	0
8	38773	0	11.6	35.5	0	0
9	38773	16	9.0	29.0	1	0
10	38773	16	14.4	31.0	0	0
11	38773	16	13.4	32.0	0	0
12	38773	16	13.2	30.0	0	0
13	38773	16	14.2	30.5	0	1
14	38773	16	14.0	33.0	0	0
15	38773	16	13.2	30.2	0	0
16	38773	16	14.9	30.4	0	0
17	38773	32	12.0	26.5	2	4
18	38773	32	11.6	25.4	1	1
19	38773	32	9.4	22.0	1	3
20	38773	32	11.2	23.0	2	4
21	38773	32	13.8	25.8	1	2
22	38773	32	12.4	23.0	1	4
23	38773	32	10.2	19.0	2	4
24	38773	32	13.8	23.2	2	3
26	38773	64	11.8	20.0	2	5
27	38773	64	13.0	22.0	2	5
28	38773	64	12.2	18.0	3	5
29	38773	64	15.8	23.0	2	5
30	38773	64	10.4	17.5	2	5
32	38773	64	15.0	18.5	2	5
33	38773	64	13.8	21.8	2	5
34	38773	64	13.6	19.0	3	5
35	38773	64	10.8	16.0	3	5
36	Wild type	0	21.0	40.6	0	0
37	Wild type	0	16.0	38.0	0	0
38	Wild type	0	15.0	34.6	0	0
39	Wild type	0	17.6	32.2	0	0
40	Wild type	0	15.0	31.6	0	0
41	Wild type	0	14.0	32.0	0	0
42	Wild type	16	10.0	11.8	3	5
43	Wild type	16	8.0	10.0	3	5
44	Wild type	16	8.6	11.0	3	5

45	Wild type	16	8.0	14.0	3	5
46	Wild type	16	9.8	11.0	3	5
47	Wild type	16	10.4	14.0	3	5
48	Wild type	32	10.8	13.2	3	5
49	Wild type	32	9.0	13.0	3	5
50	Wild type	32	8.0	10.2	3	5
51	Wild type	32	11.0	14.0	4	5
52	Wild type	32	9.8	13.0	3	5
53	Wild type	32	8.0	10.8	4	5
54	Wild type	64	7.5	8.6	4	5
55	Wild type	64	11.2	12.5	4	5
56	Wild type	64	10.2	12.8	4	5
57	Wild type	64	11.5	13.0	4	5
58	Wild type	64	13.0	15.0	4	5
59	Wild type	64	9.8	11.2	4	5

Vegetative injuries:

0=normal plant

1=slight chlorosis of new leaves and stunting

2=severe chlorosis of new leaves, malformation of new leaves, and severe stunting

3=dying plant

4=dead plant

Fertility ratings:

0 = Fertile, no delay in maturity, lots of seed

1 = Some abortion, slight delay in seed set, seed

2 = Significant abortion, significant delay in seed set, some seed

3 = Very severe abortion, immature seed pots, a few seed

4 = malformed flowers; if flowered, extreme delay in flowering and no seed produced

5=dead plant

In addition, other transplastomic lines were analyzed for tolerance to spraying with various levels of glyphosate as described above. Specific activity is measured as the amount of exogenously added Phosphoenol pyruvate (PEP) converted to Shikimate-3-phosphate (S3P) per unit protein in the plant extract. Addition of glyphosate tests sensitivity of the EPSPS enzyme to glyphosate. The results are summarized in Table 7.

**Table 7**

LINE	% TOTAL	SPECIFIC	SPECIFIC	Vegetative	Reproductive
------	---------	----------	----------	------------	--------------

	SOLUBLE PROTEIN	ACTIVITY (nmol/min/mg) No gly	ACTIVITY (nmol/min/mg) +1 mM gly	tolerance (oz/acre)	tolerance (oz/acre)
Wild-type		3.4	0	0	0
pMON10154	0.04	19.0	18.4	128	64
pMON45201	-	301.6	221.2	32	32
pMON45204	-	339.9	371.8	128	64
pMON30123	0.001	4.0	0	0	0
pMON30130	0.002	6.2	0	0	0
pMON38773	0.2	16.7	6.7	32	16
pMON38798	0.2	17.2	14.7	32	16
pMON45259	>12.0	-	-	128	64
pMON49218	>12.0	-	-	128	64

These data demonstrate that high levels of glyphosate tolerance can be obtained in transplastomic plants expressing various EPSPS sequences. In particular, lines pMON45204, pMON45259, and pMON49218 provide tolerance to glyphosate applied at levels of at least 128 oz/acre on vegetative tissues, and at least 64 oz/acre on reproductive  
5 tissues.

Furthermore, constructs pMON42259 and pMON49218 provide for high level expression of CP4 EPSPS from plant plastids transformed with these constructs. In particular, expression levels of greater than about 12 percent total soluble protein are obtained in constructs employing sequences encoding the first 14 amino acids of GFP  
10 fused to the N-terminus of CP4.

### 3F. BT/BXN Analysis

Homoplasmic tobacco plants of the lines 5175 and 5197 are sprayed with Buctril herbicide at a concentration of 4% to test for whole plant tolerance.

Results of the spray test with Buctril demonstrated that all 5197 lines expressing *bxn* were completely resistant when sprayed with a solution containing 4% Buctril herbicide.

Two lines out of six 5175 lines tested were completely resistant to the herbicide when sprayed with a 4% solution containing Buctril.

### 3G. Norflurazon Resistance Analysis

An experiment was set up to determine the efficacy of the *Crt I* trait with respect to resistance to the herbicide Norflurazon. Three 5177 transformed lines, 74-1B-P, 74-2-A, and 74-7-C and three control lines were planted. Plants were grown for seven weeks and then watered with a 3 $\mu$ M Norflurazon solution. Plants negative for the presence of the *crtI* plastid-borne gene were bleached by Norflurazon treatment, positive plants stayed green and continued to grow.

The results show that the three homoplasmic 5177 tobacco lines were resistant to the 3 $\mu$ M Norflurazon solution, while the control plants were all susceptible to the solution (Table 8).

Table 8

Line	Control/Transgenic	Result
Xanthi	Control	Susceptible
2560A Xanthi	Control	Susceptible
75-5D-A	Control	Susceptible
74-1B-P	homoplasmic	Resistant
74-2-A	homoplasmic	Resistant
74-7-C	homoplasmic	Resistant

## 20 Example 4 Analysis of hGH Transgenic Tobacco Plants

### 4A. Southern Analysis

Transformed plants selected for *aadA* marker gene expression are analyzed to determine whether the entire plastid content of the plant has been transformed (homoplastic transformants). Homoplasmic plants are selected using Southern hybridization for further cultivation.

5            Genomic DNA is isolated from transformed tobacco plants, electrophoresed, and transferred to filters as described in Svab *et al.* ((1993), *Proc Natl Acad Sci*, 90:913-917).

             Homoplasmic tobacco plants transformed to express hGH were identified using a probe prepared from a 2.4 kb *EcoRI/EcoRV* fragment from the vector pOVZ2 (similar to pOVZ15 described in Zoubenko, *et al.* 1994, *supra*). The 2.4 kb probe fragment  
10           encompasses part of the targeting sequence.

             The complete disappearance of the 3.27 Kb native tobacco *BamHI* fragment in the lines with a probe covering the region of integration, and the appearance of the expected size band for the inserted DNA fragments in those transformants, 5.6 kb, establishes that the transformed plants are homoplasmic for the intended constructs.

15

#### 4B. Protein Expression Analysis

             Homoplasmic tobacco lines expressing hGH and nuclear tobacco transformants are used to determine the expression of the hGH protein. Western blot analysis was performed on tobacco lines containing constructs pWRG4838, pMON38755 and pMON38794 for  
20           plastid expression and an ELISA assay was used for transgenic tobacco lines containing pWRG4744 and pWRG4747 for nuclear expression of hGH.

             Total protein extractions and western blot procedures were performed as described above, with the exception of the primary antibody was raised against hGH.

**Table 9****Expression Levels of hGH in Tobacco Nuclear Genome and Plastid genome**

Construct	Expression	Expression Level % Total Soluble Protein
pWRG4744	nuclear	0.002-0.125%
pWRG4747	nuclear	0.002-0.025%
pWRG4838	plastid	0.2%
pMON38755	plastid	1.0%
pMON38794	plastid	7.0%

Results of the Western analysis (Table 9) demonstrates that hGH expressed in

5 plastids of plant cells accumulates to significantly higher levels than hGH expressed in the nucleus and targeted to either the cytoplasm or plastid of plant cells. Tobacco plants transformed to express hGH in the nucleus accumulated hGH levels of 0.002% (cytoplasmic targeted) to 0.025% (plastid targeted) of total soluble leaf protein, while tobacco plants expressing hGH in the plastid accumulated hGH levels of 0.2% to 7.0% of

10 the total soluble leaf protein as hGH. Furthermore, homoplasmic tobacco plants expressing hGH directed from the *Prrn*/G10L regulatory sequences accumulate 35 fold higher levels of hGH than homoplasmic tobacco plants expressing hGH directed from the *PpsbA* promoter sequence.

15 **4C. Characterization of hGH Protein Expressed in the Plastid**

In order to determine whether the hGH expressed from plastids was properly processed, experiments were performed to determine correct folding and bioactivity.

Two bottom leaves of transplastomic tobacco lines containing pMON38794 were used to extract and purify hGH. Large veins were removed from the excised leaves, and

20 the leaf tissue was cut into small sections (approximately 0.5 cm<sup>2</sup>). The leaf pieces were flash frozen in liquid nitrogen and ground to a fine powder in a chilled mortar and pestle.

Ten grams of frozen, ground leaf tissue was added to ice cold 100 mM Tris base solution (30 ml) and mixed vigorously by vortexing for 5 minutes. The solution was filtered through a single layer of cheese cloth.

From the filtered solution, three separate samples were prepared. The first sample was prepared by centrifuging 4 ml of the filtrate for 1 minute at 16,000 rpm. The centrifugate was aliquoted into 1 ml vials and frozen in dry ice. The remaining filtrate was centrifuged for 10 minutes at 4800 rpm, and several 0.5 ml aliquots were frozen as above for the second sample. To the remaining centrifuged filtrate (approximately 25 ml), 200  $\mu$ l of glacial acetic acid was added to lower the pH from 8.2 to 4.56. The solution was centrifuged at 4800 rpm for 30 minutes, and the supernatant was frozen over dry ice for the third sample.

Total soluble protein (TSP, Table 10) was calculated in these samples by standard protein assay procedures (Maniatis, ), and the percent purity of hGH was calculated based on results from Western blot analysis using known concentrations of starting material.

Table 10

Sample ID	TSP mg/mL	GP2000 mg/L	% Purity
Filtered Extract immediately centrifuged and frozen	6.3	28	0.45 %
Filtered extract centrifuged at 4800 rpm for 10 min and frozen	6.4	28	0.45
pH adjusted and centrifuged extract	0.75	21	2.8%

The pH adjusted and centrifuged extract was purified by Reverse Phase-HPLC (RP-HPLC) for electrospray mass spectrometry and amino-terminal amino acid sequencing. RP-HPLC was performed using a Perkin-Elmer series 200 pump and autosampler and a Vydac C8 (250 by 4.6 mm) RP-HPLC column. 750 microliters of sample was loaded onto the column equilibrated with 20 mM trifluoroacetic acid (TFA) and 50 % acetonitrile. After loading, the column was washed for 2 minutes with 50 %



5 The results of the RP-HPLC analysis are shown in figure 3. Peak I (tallest peak) has the retention time expected for properly folded, native 22 kDa GP2000. This peak was collected and dried down in a Savant Speed-Vac for amino terminal sequencing and electrospray mass spectrometry.

The results of the mass spectrometry of the RP-HPLC peak I shows 4 major protein species of different molecular mass. The 21,997 kDa species represents the predicted mass of hGH with the predicted N-terminal Phe removed by over-cleavage of the Ubiquitin protease. The 22,124 kDa species represents the predicted mass of properly processed, correct amino acid sequence of hGH. The 22,507 kDa and 22,664 kDa species are thought to represent an hGH with the N-terminal Phe and hGH which have been modified during plant extraction procedures, respectively. The calculated molecular mass of the proteins suggests that the hGH expressed from the plastid is properly folded (i.e. the correct disulfide bonds are created).

PNSD/CID JWO      MMR3022A.3 1A>

#### 4D. Bioactivity of hGH Expressed in Plant Plastids

Bioactivity of the pH adjusted and centrifuged extract was tested using cells from an Nb2 cell line. These cells proliferate in the presence of growth hormone and other estrogenic type compounds. The assay involves putting various concentrations of growth hormone-containing extract into a 96 well plate. Then a constant amount of cells are added to each well. The plate is incubated for 48hrs and then a reagent called MTS is added. Metabolizing cells take up the MTS and convert it to a blue colored substance. The more cells there are the more blue color in the well. The blue color is measured using a spectrophotometer. The number of cells should be proportional to the concentration of growth hormone in the media. At some high concentration one expects that the cells will become saturated with growth hormone and that the dose response will level off. At very low hGH concentrations essentially no enhanced growth is seen. A sigmoidal shape graph is expected to be produced graphing the cell number (or absorbance) vs hGH concentration graph.

The results of the bioactivity assay (Figure 5) demonstrates that the hGH expressed from a plant plastid has a sigmoidal shape when graphed as absorbance vs hGH concentration.

#### Example 5 Analysis of Aprotinin Transplastomic Tobacco Plants

20

##### 5A. Western Analysis of Aprotinin Expression in Plastids

Homoplasmic tobacco lines expressing are used to determine the expression of the aprotinin protein. Western blot analysis was performed on tobacco lines containing constructs pCGN6146, pCGN6147, pCGN6154 and pCGN6156 for plastid expression of aprotinin.

Total protein extractions and western blot procedures were performed as described above, with the exception of the primary antibody was raised against aprotinin.

The results of the Western analysis indicate that aprotinin is expressed from the T7 polymerase promoter when the aprotinin coding sequence is fused with either the PetA or

full length GUS gene. Furthermore, these results indicate that the petA sequence efficiently targets the aprotinin protein to the plant cell thylakoid.

5           All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

10           Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claim.

## CLAIMS

What is claimed is:

- 5
1. A construct comprising the following components in the 5' to 3' direction of transcription:
- 10
- a) a promoter functional in a plant plastid;
  - b) a DNA sequence which is capable of conferring tolerance in a plant cell to at least one herbicide compound when said DNA sequence is transcribed in plastids of said plant cell; and
  - c) a transcription termination region.
- 15
2. The construct according to Claim 1, wherein said construct further comprises (d) a gene encoding a selectable marker for selection of plant cells comprising a plastid expressing said marker and (e) DNA regions of homology to the genome of said plastid, wherein said regions of homology in (e) flank components (a), (b), (c) and (d).
- 20
3. The construct according to Claim 1, wherein said construct further comprises (f) a ribosome binding site joined to said promoter component (a).
- 25
4. The construct according to Claim 3, wherein said ribosome binding site (f) is from a leader sequence selected from the group consisting of sites derived from plastid, bacterial or bacteriophage leader sequences.
- 30
5. The construct according to Claim 4, wherein said ribosome binding site is selected from the group consisting of the binding site of the gene 10 leader and the rbcLRBS site.
- 35
6. The construct according to Claim 1 wherein said DNA sequence encodes a gene which confers tolerance to the herbicide glyphosate.
- 40
7. The construct according to Claim 6 wherein said DNA encoding sequence encodes a glyphosate-tolerant 5-enolpyruvylshikimate-3-phosphate synthase.
- 45
8. The construct according to Claim 7 wherein said DNA encoding sequence is selected from the group consisting of the *E. coli* or *Salmonella aroA* gene, the CP4 gene, mutant petunia EPSPS, mutant EPSPS gene of *Psuedomonas* strain LBAA, and the *Bacillus subtilis aroE* gene.
9. The construct according to Claim 6 wherein said DNA encoding sequence encodes a glyphosate-modifying enzyme.
10. The construct according to Claim 9 wherein said glyphosate-modifying enzyme is selected from the group consisting of the *gox*, *hph*, *glpA* and *glpB*.

11. The construct according to Claim 1 wherein said DNA encoding sequence is the native encoding sequence to said gene.

5 12. The construct according to Claim 1 wherein said DNA encoding sequence is a synthetic encoding sequence to said gene.

13. The construct according to Claim 1 wherein said DNA sequence encodes a sulphonylurea-tolerant AHAS gene.

10 14. The construct according to Claim 1 wherein said DNA sequence encodes a imidizalinone-tolerant AHAS gene.

15 15. The construct according to Claim 14 wherein said DNA encoding sequence is the ALS gene.

16. The construct according to Claim 1 wherein said DNA sequence encodes a phosphinothricin-tolerant gene.

20 17. The construct according to Claim 16 wherein said DNA encoding sequence is the BAR gene.

18. The construct according to Claim 1 wherein said DNA sequence encodes an enzyme of the carotenoid pathway.

25 19. The construct according to Claim 18 wherein said DNA encoding sequence is the crtI gene.

30 20. The construct according to Claim 1 wherein said DNA sequence encodes a bromoxynil-tolerant gene.

21. The construct according to Claim 20 wherein said bromoxynil-tolerant gene is the BXN gene.

35 22. A plant cell plastid containing the construct according to Claim 1.

23. A plant, plant seed, plant cell or progeny thereof containing a plant plastid according to Claim 22.

40 24. A method for producing tolerance of a herbicide in a plant cell, wherein said method comprises transforming plastids of said plant cell with a construct comprising the following as operably joined components in the 5' to 3' direction of transcription:

45 a) a promoter functional in a plant plastid;  
b) a DNA sequence which is capable of conferring in a plant cell tolerance to at least one herbicide compound when said DNA sequence is transcribed in plastids of said plant cell; and

c) a transcription termination region,  
and growing plant cells comprising said transformed plastids under conditions wherein said DNA sequence is transcribed whereby plant cells containing said plant plastids are rendered tolerant to applications of said at least one herbicide compound.

25. The method according to Claim 24, wherein said construct further comprises (d) a gene encoding a selectable marker for selection of plant cells comprising a plastid expressing said marker and (e) DNA regions of homology to the genome of said plastid, wherein said regions of homology in (e) flank components (a), (b), (c) and (d).

26. The method according to Claim 24, wherein said construct further comprises a second DNA sequence of interest.

27. The method according to Claim 26, wherein said second DNA sequence comprises a gene expressed from an promoter independent from said promoter (a).

28. The method according to Claim 26, wherein said second DNA sequence comprises a gene expressed from said promoter (a) as a polycistronic message with said DNA sequence in (b).

29. The method according to Claim 26, wherein said second DNA sequence comprises a gene other than a gene capable of conferring in a plant cell tolerance to said herbicide compound.

30. The method according to Claim 26, wherein said second DNA sequence comprises a gene capable of conferring in a plant cell tolerance to a second herbicide compound.

31. The method according to Claim 24, wherein said construct further comprises (f) a ribosome binding site joined to said promoter component (a).

32. The construct according to Claim 31, wherein said ribosome binding site is selected from the group consisting of the binding site of the gene 10 leader and the rbcLRBS site.

33. The method according to Claim 24 wherein said DNA sequence encodes a gene which confers tolerance to the herbicide glyphosate.

34. The method according to Claim 33 wherein said DNA encoding sequence encodes a glyphosate-tolerant 5-enolpyruvylshikimate-3-phosphate synthase.

35. The method according to Claim 34 wherein said DNA encoding sequence is selected from the group consisting of the E. coli or Salmonella aroA gene, the CP4 gene, mutant petunia EPSPS, mutant EPSPS gene of *Psuedomonas* strain LBAA and the *Bacillus subtilis* aroE gene.

36. The method according to Claim 33 wherein said DNA encoding sequence encodes a glyphosate-modifying enzyme.

37. The method according to Claim 36 wherein said glyphosate-modifying enzyme is selected from the group consisting of the gox, hph, glpA and glpB.

38. The method according to Claim 24 wherein said DNA encoding sequence is the native encoding sequence to said gene.

5 39. The method according to Claim 24 wherein said DNA encoding sequence is a synthetic encoding sequence to said gene.

40. The method according to Claim 24 wherein said DNA sequence encodes a sulphonylurea-tolerant AHAS gene.

10 41. The method according to Claim 24 wherein said DNA sequence encodes a imidizalinone-tolerant AHAS gene.

42. The method according to Claim 41 wherein said DNA encoding sequence is the ALS gene.

15 43. The method according to Claim 24 wherein said DNA sequence encodes an enzyme of the carotenoid pathway.

44. The method according to Claim 43 wherein said DNA encoding sequence is the crtI gene.

45. The method according to Claim 24 wherein said DNA sequence encodes a bromoxynil-tolerant gene.

25 46. The method according to Claim 45 wherein said bromoxynil-tolerant gene is the BXN gene.

47. An herbicide tolerant plant cell produced according to the method of Claim 24.

30 48. A plant, plant seed or plant part comprising a plant cell according to Claim 47.

49. A plant cell according to Claim 47 and comprising greater than about 0.01 % of total soluble protein as a protein expressed from said herbicide-tolerance gene.

35 50. A plant cell according to Claim 47 and comprising greater than about 0.1 % of total soluble protein as a protein expressed from said herbicide-tolerance gene.

51. A plant cell according to Claim 47 and comprising greater than about 0.2 % of total soluble protein as a protein expressed from said herbicide-tolerance gene.

40 52. A plant cell according to Claim 47 and comprising 1 % or more of total soluble protein as a protein expressed from said herbicide-tolerance gene.

53. A plant cell according to Claim 47 and comprising 12 % or more of total soluble protein as a protein expressed from said herbicide-tolerance gene.

45 54. A plant cell according to Claim 47 wherein said herbicide-tolerance gene is a glyphosate-tolerant 5-enolpyruvylshikimate-3-phosphate synthase.

55. A plant, plant seed or plant part comprising a plant cell according to Claim 53.

56. A plant according to method of Claim 55 tolerant of the herbicide glyphosate when said herbicide is applied at a rate of about 16 ounces or greater per acre.

5

57. A plant according to method of Claim 55 tolerant of the herbicide glyphosate when said herbicide is applied at a rate of about 32 ounces or greater per acre.

57. A plant according to method of Claim 55 tolerant of the herbicide glyphosate when said herbicide is applied at a rate of about 64 ounces or greater per acre.

10

58. A plant according to method of Claim 55 tolerant of the herbicide glyphosate when said herbicide is applied at a rate of about 128 ounces or greater per acre.

60. A method wherein herbicide-tolerance produced according to the method of Claim 24 is used to select cells transformed by said construct from non-transformed cells.

15

61. A method according to Claim 60 wherein said herbicide-tolerance gene confers tolerance to the herbicide glyphosate.

20

62. A method according to Claim 61 wherein selection for said herbicide tolerance is made on media containing glyphosate at a concentration from at least about 50  $\mu$ M to about 200  $\mu$ M.

63. A method according to Claim 62 wherein selection for said herbicide tolerance is made on media containing glyphosate at a concentration from about 1 mM.

25

64. A plant cell produced according to the method of Claim 60.

65. The method according to Claim 24, wherein said plant cells are tolerant to applications of herbicide amounts selected from the group consisting of 16 ounces/acre, 32 ounces/acre, 64 ounces/acre, and 128 ounces/acre.

30

66. The method according to Claim 65, wherein said herbicide is glyphosate.



5'-AAT TGT AGA AAT AAT TTT GTT TAA CTT TAA GAA GGA GAT ATA CC-3'  
TTA ACA TCT TTA TTA AAA CAA ATT GAA ATT CTT CCT CTA TAT GG

**FIGURE 1**

Arg-Pro-Asp-Phe-Cys-Leu-Glu-Pro-Pro-Tyr-Thr-Gly-Pro-Cys-  
Lys-Ala-Arg-Ile-Ile-Arg-Tyr-Phe-Tyr-Asn-Ala-Lys-Ala-Gly-  
Leu-Cys-Gln-Thr-Phe-Val-Tyr-Gly-Gly-Cys-Arg-Ala-Lys-Arg-  
Asn-Asn-Phe-Lys-Ser-Ala-Glu-Asp-Cys-Met-Arg-Thr-Cys-Gly-  
Gly-Ala

Figure 2

3 / 7




Figure 3




Figure 4

Summary Graph




Figure 5

	EcoRI	SacI	KpnI	Prrn	
1	GAATTCGAGC	TCGGTACCCA	AAGCTCCCC	GCCGTCGTT	AATGAGAATG
51	GATAAGAGGC	TCGTGGGATT	<u>GACGTGAGGG</u>	GGCAGGGATG	<u>GCTATATTTC</u>
			-35	-10	
	PEP	G10L			
101	TGGGAGCGAA	CTCCGGGCGA	ATTGTAGAA	TAATTTTGT	TAACTTTAAG
	RBS	NcoI			
151	AAGGAGATAT	ACCCATGG			

Figure 6

EcorI KpnI Prrn  
 1 GAATTCGGTA CCCCCGTCGT TCAATGAGAA TGGATAAGAG GCTCGTGGGA  
 -35  
 51 TTGACGTGAG GGGCAGGGA TGGCTATATT TCTGGGAGCG AACTCCGGGC  
 -10  
 PEP  
 101 GAATACTGAA GCGCTTGGAT ACAAGTTATC CTGGGAAGGA AAGACAATTC  
 NEP  
 G10L  
 151 CGGATCCTCT AGAAATAATT TTGTTTAACT TTAAGAAGGA GATATACCC ATG  
 Met  
 14aa GFP  
 GGT AAA GGA GAA GAA CTT TTC ACT GGA GTT GTC CCA  
 Gly Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro  
 AGC ATG  
 Ser Met

Figure 7

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/15472

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N15/55 C12N15/60 C12N15/82 C12N5/10  
A01H5/00 A01H5/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 32011 A (CIBA GEIGY AG ;VOLRATH SANDRA L (US); JOHNSON MARIE A (US); POTTER) 4 September 1997 (1997-09-04)  see the whole document; esp. examples 34-38	1-6, 9-27, 29-33, 36-48, 60-64
X	US 5 693 507 A (DANIELL HENRY ET AL) 2 December 1997 (1997-12-02)  see the whole document; esp. example 4  -/-	1-27, 29-48, 54,60-64



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*A\* document member of the same patent family

Date of the actual completion of the international search

2 May 2000

Date of mailing of the international search report

10. 05. 2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Kania, T



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/15472

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DANIELL ET AL: "Containment of herbicide resistance through genetic engineering of the chloroplast genome" NATURE BIOTECHNOLOGY, US, NATURE PUBLISHING, vol. 16, April 1998 (1998-04), page 345-348 XP002090409 ISSN: 1087-0156 the whole document	1-27, 29-48, 54, 60-64
X	DATTA ET AL: "Transformation of the tobacco chloroplast genome with the aroA gene to confer glyphosate tolerance" PLANT PHYSIOLOGY, US, AMERICAN SOCIETY OF PLANT PHYSIOLOGISTS, ROCKVILLE, MD, vol. 111, no. 2, 1996, page 168 XP002090406 ISSN: 0032-0889 abstract	1-8, 11, 12, 22-25, 31-35, 38, 39, 47, 48, 54, 60-64
A	MCBRIDE ET AL: "Amplification of a chimeric Bacillus gene in chloroplasts leads to an extraordinary level of an insecticidal protein in tobacco" BIO/TECHNOLOGY, US, NATURE PUBLISHING CO. NEW YORK, vol. 13, 1995, page 362-365 XP002090404 ISSN: 0733-222X the whole document	1-8, 11, 12, 22-25, 31-35, 38, 39, 47-66
A	WO 98 11235 A (CIBA GEIGY AG ; HEIFETZ PETER (US); LEBEL EDOUARD (US); UKNES SCOTT) 19 March 1998 (1998-03-19)  see example C	1-8, 11, 12, 22-25, 31-35, 38, 39, 47-66
A	WO 95 24493 A (CALGENE INC ; MCBRIDE KEVIN E (US); STALKER DAVID M (US)) 14 September 1995 (1995-09-14)  the whole document	1-8, 11, 12, 22-25, 31-35, 38, 39, 47-66
A	WO 95 16783 A (CALGENE INC ; MCBRIDE KEVIN E (US); STALKER DAVID M (US)) 22 June 1995 (1995-06-22) cited in the application  the whole document	1-8, 11, 12, 22-25, 31-35, 38, 39, 47-66

-/--

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/15472

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 293 358 A (MONSANTO CO) 30 November 1988 (1988-11-30)  the whole document	1-8,11, 12, 22-25, 31-35, 38,39, 47-66
A	WO 92 04449 A (MONSANTO CO) 19 March 1992 (1992-03-19) cited in the application  the whole document	1-8,11, 12, 22-25, 31-35, 38,39, 47-66
A	WO 97 04103 A (LEBRUN MICHEL ; SAILLAND ALAIN (FR); FREYSSINET GEORGES (FR); RHONE) 6 February 1997 (1997-02-06)  the whole document	1-8,11, 12, 22-25, 31-35, 38,39, 47-66
A	US 5 589 612 A (JILKA JOSEPH M ET AL) 31 December 1996 (1996-12-31) see esp. p.10 l.9-26; p.14 l.48-60	60,61,64
A	ZHOU H ET AL: "GLYPHOSATE-TOLERANT CP4 AND GOX GENES AS A SELECTABLE MARKER IN WHEAT TRANSFORMATION" PLANT CELL REPORTS, DE, SPRINGER VERLAG, vol. 15, no. 3/04, 1 December 1995 (1995-12-01), pages 159-163, XP000578832 ISSN: 0721-7714 the whole document	61-64
A	US 5 633 437 A (BERNASCONI PAUL ET AL) 27 May 1997 (1997-05-27)  see the whole document; esp. p.5 l.14-34  -/-	1-5, 11-15, 22-27, 29-32, 38-42, 47-53, 55,65

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/15472

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MISAWA N ET AL: "FUNCTIONAL EXPRESSION OF THE ERWINIA UREDOVORA CAROTENOID BIOSYNTHESIS GENE CRTL IN TRANSGENIC PLANTS SHOWING AN INCREASE OF BETA-CAROTENE BIOSYNTHESIS ACTIVITY AND RESISTANCE TO THE BLEACHING HERBICIDE NORFLURAZON" PLANT JOURNAL, GB, BLACKWELL SCIENTIFIC PUBLICATIONS, OXFORD, vol. 4, no. 5, 1 January 1993 (1993-01-01), pages 833-840, XP002017202 ISSN: 0960-7412 the whole document	1-5,11, 12,18, 19, 22-25, 31,32, 38,39, 43,44, 47-53, 55,65
A	EP 0 337 899 A (RHONE POULENC AGROCHIMIE) 18 October 1989 (1989-10-18)  the whole document	1-5,11, 12,20, 22-25, 31,32, 38,39, 45, 47-53, 55,65
A	WO 98 20144 A (HAWKES TIMOTHY ROBERT ;KNIGHT MARY ELIZABETH (GB); THOMPSON PAUL A) 14 May 1998 (1998-05-14)  see the whole document; esp. examples 4-9	1-6, 9-27, 29-33, 36-53, 55-58, 65,66
P,X	WO 99 05265 A (BAILEY ANA MARIA ;INVEST Y DE ESTUDIOS AVANZADOS (MX); BLOWERS ALA) 4 February 1999 (1999-02-04)  the whole document	1-6, 9-12,16, 17, 22-33, 36-39, 47,48, 56,65,66

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 99/15472

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99/15472

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-6,11,12,22-25,31-33,38,39,47-53,  
55-66 partially; 7,8,34,35,54 completely

Plant plastid expression constructs comprising: a promoter functional in a plant plastid, a DNA sequence, native or synthetic, capable of conferring tolerance in the plant cell to at least one herbicide compound, and a transcription termination region. Said constructs further comprising a selectable marker as specified, DNA regions of homology to the plastid genome flanking the other component, a ribosome binding site, preferably the binding site of the gene 10 leader or rbcLRBS. Said DNA sequence encoding a gene which confers tolerance to glyphosate, preferably an EPSPS encoding gene out of a selection as claimed .  
Plant cell plastids, plant cells, plants, and seeds comprising said constructs.

A method for producing tolerance of a herbicide in a plant cell by transforming plant plastids with said constructs and growing plant cells under conditions wherein said DNA sequence is transcribed and the plant cells are rendered tolerant to herbicide applications. Said method employing the said constructs as defined above.

Plants and plant cells derived therefrom comprising an amount of said protein as claimed, and tolerant to herbicide concentrations as claimed.

A method wherein herbicide-tolerance produced according to said method is used to select transformed cells from non-transformed cells under selection conditions as defined, and plant cell produced thereby.

2. Claims: 1-6,11,12,22-25,31-33,38,39,47-53,  
55-66 partially; 9,10,36,37 completely

idem, the DNA sequence encoding a glyphosate-modifying enzyme out of a selection as claimed

3. Claims: 1-5,11,12,22-25,31,32,38,39,47-53,55,60,64,  
65 partially; 13-15,40-42 completely

idem, said DNA sequence encoding a sulphonylurea-tolerant AHAS gene, a imidizalinone-tolerant AHAS gene, preferably a ALS gene

4. Claims: 1-5,11,12,22,23 partially; 16,17 completely

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99/15472

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Plant plastid expression constructs comprising: a promoter functional in a plant plastid, a DNA sequence, native or synthetic, capable of conferring tolerance in the plant cell to at least one herbicide compound, and a transcription termination region. Said constructs further comprising a selectable marker as specified, DNA regions of homology to the plastid genome flanking the other component, a ribosome binding site, preferably the binding site of the gene 10 leader or rbcLRBS. Said DNA sequence encoding a phosphinotricin-tolerant gene, preferably the BAR gene. Plant cell plastids, plant cells, plants, and seeds comprising said constructs.

5. Claims: 1-5,11,12,22-25,31,32,38,39,47-53,55,60,64,  
65 partially; 18,19,43,44 completely

as subject 3, the DNA sequence encoding an enzyme of the carotenoid pathway, preferably the crtI gene

6. Claims: 1-5,11,12,22-25,31,32,38,39,47-53,55,60,64,  
65 partially; 20,21,45,46 completely

idem, the DNA sequence encoding a bromoxynil-tolerant gene, preferably the BXN gene

7. Claims: 24 partially; 26-30 completely

A method for producing tolerance of a herbicide in a plant cell by transforming plant plastids with a construct as defined and growing plant cells under conditions wherein said DNA sequence is transcribed and the plant cells are rendered tolerant to herbicide applications.

The method wherein said construct further comprises a second DNA sequence of interest. Said DNA sequence optionally being expressed independently from said promoter, or together with the first DNA sequence from the same promoter as a polycistronic message. Said DNA sequence comprising a gene other than a gene capable of conferring in a plant cell tolerance to said herbicide compound, or capable of conferring tolerance to a second herbicide compound.

### Information on patient family members

PCT/US 99/15472

Form PCT/ISA/210 (patent family annex) (July 1992)

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No  
PCT/US 99/15472

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9204449 A		EP 0546090 A ES 2089232 T GR 3021058 T JP 6500472 T US 5633435 A US 5627061 A US 5804425 A	16-06-1993 01-10-1996 31-12-1996 20-01-1994 27-05-1997 06-05-1997 08-09-1998
WO 9704103 A	06-02-1997	FR 2736926 A AU 6619196 A BG 102247 A BR 9609792 A CA 2223875 A CN 1196088 A CZ 9800174 A EP 0837944 A HU 9900463 A JP 11510043 T PL 324572 A SK 6498 A	24-01-1997 18-02-1997 30-04-1999 16-03-1999 06-02-1997 14-10-1998 17-06-1998 29-04-1998 28-05-1999 07-09-1999 08-06-1998 04-11-1998
US 5589612 A	31-12-1996	NONE	
US 5633437 A	27-05-1997	NONE	
EP 0337899 A	18-10-1989	FR 2629098 A AT 188250 T AU 629539 B AU 3159189 A BR 8901499 A CA 1314505 A CN 1036038 A, B DD 294040 A DE 68929125 D DK 144689 A EG 20026 A EP 0894865 A HU 52158 A, B IL 89614 A JP 2009374 A KR 145703 B NZ 228458 A PL 278448 A RO 107001 A TR 23682 A US 5559024 A US 5773698 A ZA 8902146 A	29-09-1989 15-01-2000 08-10-1992 28-09-1989 14-11-1989 16-03-1993 04-10-1989 19-09-1991 03-02-2000 24-09-1989 30-10-1997 03-02-1999 28-06-1990 19-01-1996 12-01-1990 01-08-1998 25-06-1991 27-12-1989 30-08-1993 06-06-1990 24-09-1996 30-06-1998 29-11-1989
WO 9820144 A	14-05-1998	AU 4789597 A BG 103462 A BR 9712695 A CN 1236394 A CZ 9901616 A EP 0946737 A PL 333131 A	29-05-1998 31-01-2000 19-10-1999 24-11-1999 11-08-1999 06-10-1999 22-11-1999
WO 9905265 A	04-02-1999	AU 8583798 A	16-02-1999